

# Gut microbiota helps identify clinical subtypes of Parkinson's disease

Jing-Yi Wang<sup>1†</sup>, Rui Xie<sup>1†</sup>, Yun Feng<sup>2†</sup>, Min-Na Zhang<sup>1</sup>, Le He<sup>1</sup>, Bo Yang<sup>1</sup>, Hong-Gang Wang<sup>1\*</sup> and Xiao-Zhong Yang<sup>1\*</sup>

**Key words** Parkinson's disease, Gut microbiota, Functional magnetic resonance imaging

Dear Editor,

The role of the microbiota-gut-brain axis in Parkinson's disease (PD) has garnered increasing attention[1]. Among the various empirical subtype systems of PD, motor subtype classification into tremor-dominant (TD) and postural instability and gait difficulty (PIGD) is the most commonly utilized[2]. However, the relationship between gut microbiota and clinical subtypes of PD remains unclear.

This study aimed to investigate the distinctions in gut microbiota between the two motor subtypes by obtaining fresh stool samples from patients with PD for metagenomic sequencing, along with assessing brain function using resting-state functional magnetic resonance imaging (rs-fMRI). Subsequently, differences in brain mean amplitude of low-frequency fluctuation (mALFF) values and their association with gut microbiota were analyzed.

A total of 24 patients with PD were included in the study, comprising 12 PIGD, 9 TD, and 3 indeterminate (IND) subtype patients (Additional file 1: Materials and methods). Following exclusion of the 3 IND subtype patients, no significant differences in age, sex, body mass index (BMI), Hoehn-Yahr score, Unified Parkinson's Disease Rating Scale part III (UPDRS-III) score, Hamilton Depression Scale (HAMD) score, Hamilton Anxiety Scale (HAMA) score, drug use, duration of disease, and medical history including hypertension, diabetes, heart disease, and cerebral infarction were observed between the TD and PIGD groups ( $P>0.05$ ; Additional file 1: Table S1). Alpha diversity of the gut microbiota was evaluated using Chao1, Shannon, and Simpson indices, while beta diversity was assessed through principal coordinates analysis (PCoA) and non-metric multi-

dimensional scaling (NMDS). However, the analysis revealed no significant variations in alpha and beta diversities between the two groups ( $P>0.05$ ; Additional file 1: Fig. S1).

The gut microbiota was further analyzed at various taxonomic levels using a cladogram and bar plot generated by Linear discriminant analysis effect size (LEfSe). The PIGD group exhibited enrichment in *Bacteroidetes*, *Bacteroidia*, and *Bacteroidales*, while the TD group showed predominant enrichment for *Firmicutes*, *Clostridiaceae*, and *Clostridium* (Fig. 1a). At the phylum level, the proportion of *Bacteroidetes* was significantly higher in PIGD group compared with TD group (45.57% vs. 23.64%,  $P=0.003$ ; Fig. 1b).

A binary logistic regression model was developed using the proportion of *Bacteroidetes* to predict the motor subtypes of patients with PD. The ROC curve analysis yielded an AUC of 0.8889 ( $P=0.0028$ ), with a cut-off value of 23.04%, sensitivity of 75.75%, and specificity of 66.67% (Fig. 1c). Additionally, at the genus level, PIGD group exhibited significantly lower abundance of *Clostridium*, *Ruminococcus*, *Coprococcus*, *Dorea*, *Enterobacter*, *Anaerotignum*, *Tyzzerella*, *Bacillus*, and *Evtetpia* compared with TD group ( $P<0.05$ ; Additional file 1: Fig. S2).

Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis was conducted to explore differences in gene abundance between TD and PIGD patients, revealing significant enrichment in various categories including environmental information processing, human diseases, cellular processes, genetic information processing, and metabolism. Notably, the metabolic pathway emerged as the most significantly enriched pathway in KEGG pathway classification (Additional file 1: Fig. S3).


Functional brain MRI data from these patients with PD were utilized for brain mALFF analysis in this study. In comparison with TD group, individuals in PIGD group exhibited reduced mALFF signals in the bilateral median cingulate and paracingulate gyri, precuneus, and parietal lobes

<sup>†</sup>Jing-Yi Wang, Rui Xie, and Yun Feng contributed equally to this work

\*Correspondence: jgzwhg@njmu.edu.cn; hayyyzh@njmu.edu.cn

<sup>1</sup>Department of Gastroenterology, the Affiliated Huai'an No.1 People's Hospital of Nanjing Medical University, Huai'an 223301, Jiangsu, China

Full list of author information is available at the end of the article



**Fig. 1 Microbiota-gut-brain axis in PD.**

a. Linear discriminant analysis effect size (LEfSe) analysis and Cladogram generated by LEfSe analysis; b. Different proportion of phylum and statistical difference between the TD and PiGD subtypes ( $P=0.003$ ); c. ROC curve; d. Significant brain changes between the TD and PiGD subtypes (gray arrows: bilateral median cingulate and paracingulate gyri; purple arrows: precuneus; brown arrow: parietal lobes); e. Spearman analysis was performed for correlation analysis. PD. Parkinson's disease; TD. Tremor-dominant; PiGD. Postural instability and gait difficulty; mALFF. Mean amplitude of low-frequency fluctuation; Corr. Correlation coefficient

(Fig. 1d). The right median cingulate and paracingulate gyri were identified as the key brain regions, and the local mALFF values were collected from each participant. Subsequently, Spearman's correlation analysis was conducted between the extracted mALFF values and various significantly distinct gut microbiotas (Fig. 1e). The findings revealed a positive correlation between the abundance of *Enterobacter*, *Tyzzarella*, *Burkholderia*, and *Leclercia* and the mALFF values ( $P < 0.05$ ). Conversely, the abundance of *Bacteroidetes* showed a negative correlation with the mALFF values ( $P < 0.05$ ).

Our research contributes valuable insights into the microbiota-gut-brain axis and underscores the potential relationship between gut microbiota and motor subtypes in PD. These findings lay the groundwork for future investigations into the pathogenesis of distinct motor subtypes. Despite being a small-scale study, we have innovatively established a connection between gut microbiota and brain function MRI, offering potential for future regulation of gut-directed therapies in PD.

#### Abbreviations

AUC: Area under the receiver operating characteristic curve; HAMD: Hamilton depression scale; HAMA: Hamilton anxiety scale; IND: Indeterminate; KEGG: Kyoto encyclopedia of genes and genomes; LEfSe: Linear discriminant analysis effect size; mALFF: Mean amplitude of low-frequency fluctuation; NMDS: Nonmetric multidimensional scaling; PD: Parkinson's disease; PIGD: Postural instability and gait difficulty; PCoA: Principal coordinates analysis; rs-fMRI: Resting-state functional magnetic resonance imaging; TD: Tremor-dominant; UPDRS: Unified Parkinson's disease rating scale.

#### Supplementary information

The online version contains supplementary material available at <https://doi.org/10.1186/s40779-024-00545-4>.

**Additional file 1. Materials and Methods. Table S1** Characteristics of different motor subtypes (TD and PIGD) in PD patients. **Fig. S1** Alpha diversity and beta diversity of the gut microbiota. **Fig. S2** Relative abundance of discriminative gut microbiota at the genus level ( $P < 0.05$ ). **Fig. S3** Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis.

#### Acknowledgements

We would like to thank the Department of Neurology of the Affiliated Huai'an No.1 People's Hospital of Nanjing Medical University for helping recruit participants and evaluate the questionnaire.

#### Authors' contributions

JYW and HGW designed the experiments; JYW and XZY performed the overall data analysis; MNZ and LH collected the samples; BY and YF contributed to figure generation; JYW, RX, and HGW wrote the manuscript. All authors read and approved the final manuscript.

#### Funding

This work was supported in part by the Jiangsu Provincial Geriatric Health Research Project (LK2021049), the Jiangsu Provincial Geriatric Health Research Project (LKM2023044), and the Huai'an Key Laboratory of Aging and Geriatric Syndrome Research (HAP202105).

#### Availability of data and materials

The data presented in this study are available upon reasonable request from the corresponding author.

#### Declarations

##### Ethics approval and consent to participate

This study was approved by the Medical Ethics Committee of Affiliated Huai'an No.1 People's Hospital of Nanjing Medical University (KY-2022-004-01).

##### Consent for publication

Not applicable.

##### Competing interests

The authors declare that they have no competing interests.

##### Author details

<sup>1</sup>Department of Gastroenterology, the Affiliated Huai'an No.1 People's Hospital of Nanjing Medical University, Huai'an 223301, Jiangsu, China.

<sup>2</sup>Department of Radiology, the Affiliated Huai'an No.1 People's Hospital of Nanjing Medical University, Huai'an 223301, Jiangsu, China.

#### References

1. Stopińska K, Radziwoń-Zaleska M, Domitrz I. The microbiota-gut-brain axis as a key to neuropsychiatric disorders: a mini review. *J Clin Med*. 2021;10(20):4640.
2. Mestre TA, Fereshtehnejad SM, Berg D, Bohnen NI, Dujardin K, Erro R, et al. Parkinson's disease subtypes: critical appraisal and recommendations. *J Parkinsons Dis*. 2021;11(2):395–404.

<https://doi.org/10.1186/s40779-024-00545-4>

**Cite this article as:** Wang JY, Xie R, Feng Y, Zhang MN, He L, Yang B, et al. Gut microbiota helps identify clinical subtypes of Parkinson's disease. *Mil Med Res*. 2024;11(1):42.