

REVIEW

Open Access

Personalized pancreatic cancer therapy: from the perspective of mRNA vaccine

Xing Huang^{1,2,3,4,5*}, Gang Zhang^{1,2,3,4,5†}, Tian-Yu Tang^{1,2,3,4,5}, Xiang Gao^{1,2,3,4,5} and Ting-Bo Liang^{1,2,3,4,5*}

Abstract

Pancreatic cancer is characterized by inter-tumoral and intra-tumoral heterogeneity, especially in genetic alteration and microenvironment. Conventional therapeutic strategies for pancreatic cancer usually suffer resistance, highlighting the necessity for personalized precise treatment. Cancer vaccines have become promising alternatives for pancreatic cancer treatment because of their multifaceted advantages including multiple targeting, minimal nonspecific effects, broad therapeutic window, low toxicity, and induction of persistent immunological memory. Multiple conventional vaccines based on the cells, microorganisms, exosomes, proteins, peptides, or DNA against pancreatic cancer have been developed; however, their overall efficacy remains unsatisfactory. Compared with these vaccine modalities, messenger RNA (mRNA)-based vaccines offer technical and conceptual advances in personalized precise treatment, and thus represent a potentially cutting-edge option in novel therapeutic approaches for pancreatic cancer. This review summarizes the current progress on pancreatic cancer vaccines, highlights the superiority of mRNA vaccines over other conventional vaccines, and proposes the viable tactic for designing and applying personalized mRNA vaccines for the precise treatment of pancreatic cancer.

Key words Pancreatic cancer, Precise therapy, Cancer vaccine, mRNA vaccine, Tumor antigen, Immune subtype

Background

The annual pancreatic cancer cases have doubled over the past two decades, increasing from 196,000 patients worldwide in 1990 to 441,000 in 2017[1]. According to the 2020 global cancer statistics, there were 495,773 new cases of pancreatic cancer[2]. Given the increase in life expectancy of the global population, the incidence of pancreatic cancer is expected to continue rising over the coming decades. Surgical intervention is currently the only curative option for pancreatic cancer management in the clinic. However, only 15%–20% of patients qualify for the corresponding surgery, attributed to the limited routine screening methods for detecting pancreatic cancer at an early stage[3]. Moreover, despite complete resection, local or distant recurrence of pancreatic cancer is often observed within two years after surgery[4]. Systematic chemotherapy has been the standard treatment for more than 80% of patients with locally advanced diseases or distant metastases for several decades. Even though gemcitabine plus nab-paclitaxel and FOLFIRINOX are the most recommended

chemotherapeutic regimens for metastatic pancreatic ductal adenocarcinoma (PDAC) treatment, acquired resistance against these drugs is common[5-7]. Immunotherapy, targeted therapy, and other promising treatments have also been tested in preclinical studies and clinical trials; however, almost all strategies show little significant advantage over conventional chemotherapy against pancreatic cancer, together with the prevalent therapeutic resistance[8,9]. Accordingly, the overall 5-year survival of pancreatic cancer patients is only about 10%, making the tumor is one of the leading causes of cancer-related mortality[10]. Obviously, there is an urgent need for highly effective alternatives for pancreatic cancer treatment.

Accumulating evidence indicates that the therapeutic resistance in pancreatic cancer is associated with its inter-tumoral and intra-tumoral heterogeneity, particularly as regards the genetic alteration and immune microenvironment[11-13]. For instance, SMAD4 mutation occurs in about 50% of PDAC patients[14], and this mutation promotes radiotherapeutic resistance by increasing the production of reactive oxygen species and inducing autophagy[15]. Appropriately 6% of pancreatic cancers display BRCA1/2 or PALB2 mutations[16,17], and the lack of mutations in these genes is associated with resistance to platinum-based chemotherapy[17,18]. Tumors without BRCA1/2

*Xing Huang and Gang Zhang contributed equally to this work.

†Correspondence: huangxing66@zju.edu.cn; liangtingbo@zju.edu.cn

²Zhejiang Provincial Key Laboratory of Pancreatic Disease, the First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310009, China

³Department of Hepatobiliary and Pancreatic Surgery, the First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China

Full list of author information is available at the end of the article

mutations are also susceptible to generating PARP inhibitor resistance[19-21]. In addition, the inter-tumoral heterogeneity in the immune microenvironment promotes resistance to immunotherapy[11]. Taking the programmed cell death 1/programmed cell death ligand 1 (PD-1/PD-L1) blockade as an example, its therapeutic efficacy is associated with the pre-infiltration of T cells[22]. Only <1% of PDAC patients with high microsatellite instability that was detected the presence of neoantigen-specific T cell immunity in tumor respond to PD-1 inhibition[23-26]. In contrast, most tumors are characterized by low immunogenicity and lack of T cell infiltration and are thus resistant to immunotherapy targeting PD-1/PD-L1. Notably, pancreatic cancer is classified into distinct subtypes based on gene expression or immune characteristics[27-29]. For instance, Moffitt *et al.*[28] identified two stromal subtypes (normal and activated) and two tumor subtypes (basal-like and classical) based on gene expression profiles. Compared with classical subtype tumors, basal-like subtype tumors exhibit a superior response to adjuvant chemotherapy. Apart from inter-tumoral heterogeneity, increasing evidence has uncovered the intratumoral heterogeneity in pancreatic cancer[30]. At least three types of intra-tumoral genetic heterogeneity have been proposed[13]. Type-1 includes mutations distinguishing tumor cells within the same primary lesions, type-2 includes mutations distinguishing tumor cells within the same metastatic lesions, while type-3 includes mutations distinguishing tumor cells among different metastatic lesions. The intra-tumoral heterogeneity largely promotes adaptive resistance to cancer therapy. Single-cell sequencing for pancreatic cancer has revealed the existence of both basal-like and classical subtypes in the same tumor, partially explaining the adaptive resistance to chemotherapy[31]. These reports highlight the heterogeneity-induced therapeutic resistance, underlining the significance of developing personalized precise treatment against pancreatic cancer. Compared with the traditional monoclonal antibodies and small molecule inhibitors, cancer vaccines offer several advantages, including minimal nonspecific effects, broad therapeutic window, low toxicity, and induction of persistent immunological memory[32,33]. Moreover, cancer vaccines can achieve precise targeting based on the characteristics in individual tumors. Therefore, vaccination is a potential approach for personalized pancreatic cancer treatment, overcoming the challenges posed by tumor heterogeneity.

Messenger RNA (mRNA) vaccine has recently become one of the most potent vaccine types in prevention and treatment of multiple diseases. The successful development of mRNA vaccine is attributed to decades of relentless and intensive

research. mRNA was discovered in 1961 and isolated for *in vitro* protein expression in 1969[34,35]. Until 1990, *in vitro* transcribed mRNA was validated able to be template to produce proteins in mouse skeletal muscle cells *in vivo*[36]. This was the first successful attempt for *in vivo* mRNA expression, setting the stage for mRNA vaccine development. Later in 1992, mRNA for vasopressin was injected and expressed in the hypothalamus, inducing physiological responses[37]. Thereafter in 1993 and 1995, mRNA was reported to induce both cellular and humoral immunity[38-40]. However, these promising findings did not attract substantial investment in the development of mRNA vaccines largely due to the perceived mRNA instability, inefficient *in vivo* delivery, and potential innate immunogenicity. Given the safety, simple design, and ease of manufacturing, research on mRNA continued. The technological advances in the modification and delivery of mRNA largely addressed these concerns. For instance, the application of modified nucleosides prevents mRNA recognition by pattern recognition receptors (PRRs), enhancing the translational efficacy[41]. Application of vehicles (e.g., lipid nanoparticle, polyplexes, and polymeric nanoparticles) promotes the *in vivo* delivery of mRNA[42]. The improvement of *in vivo* translational efficiency and delivery enhances the chance of clinical application of mRNA vaccines. The first application of personalized mRNA vaccine in humans was reported in 2017 against melanoma, and the vaccination induced specific immune activation, decreased the metastatic rate and prolonged the progression-free survival of patients[43]. In addition, multiple clinical trials on the efficacy of mRNA-based vaccination against human immunodeficiency virus have been completed[44]. Since its outbreak in 2019, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the cause of the coronavirus disease 2019 (COVID-19), has infected and caused millions of deaths globally[45,46]. Due to the threat posed by SARS-CoV-2, various treatments were rapidly developed to contain the spread of the virus. Owing to the convenience in mass production and advances in modification techniques of mRNA, two mRNA vaccines, BNT162b2 and mRNA-1273, obtained the authorization of emergency use for preventing COVID-19[47-50]. Both achieved a protective efficacy of over 90%, and were officially approved for mass vaccination by the Food and Drug Administration against SARS-CoV-2[49-53]. Of note, SARS-CoV-2 underwent multiple mutations, compromising the protective efficacy of BNT162b2 and mRNA-1273[54-56]. However, this concern has currently been solved to a large extent by updating vaccine-encoded antigens accordingly, together with optimizing the administration of

vaccination and boosting. The strategies used for overcoming SARS-CoV-2 variations provide valuable experience for developing and applying the personalized anti-pancreatic cancer mRNA vaccine. Together, the application of mRNA vaccine in other diseases lays a foundation for the development of personalized antipancreatic cancer mRNA vaccine.

This review summarizes the current advances and status of pancreatic cancer vaccines, emphasizes the superiority of mRNA-based vaccines in cancer precision treatment, and

highlights the strategy for developing personalized mRNA vaccines against pancreatic cancer.

Conventional pancreatic cancer vaccines

To date, multiple conventional vaccines against pancreatic cancer, including cell-based, microorganism-based, exosome-based, protein-based, peptide-based, and DNA-based forms (Fig. 1), are under development (completed clinical trials are summarized in Table 1, and ongoing clinical trials in Table 2).




Fig. 1 Classification of existing pancreatic cancer vaccines.

Multiple pancreatic cancer vaccines have been developed to date, including cell-based vaccines, microorganism-based vaccines, exosome-based vaccines, protein-based vaccines, peptide-based vaccines, and DNA-based vaccines. DC. Dendritic cell

Cell-based pancreatic cancer vaccines

The currently available cell-based pancreatic cancer vaccines include dendritic cell (DC)-based and tumor cell-based forms. As the most potent antigen-presenting cell (APC), DCs are usually loaded with an antigen and re-infused into patients[57]. In a phase I/II clinical trial, 12 patients with resected pancreatic and biliary cancer received mucin 1 (MUC1) peptide-loaded DC vaccine[57,58]. Four of them survived more than 4 years after the vaccination and showed no signs of recurrence. In a related study, Wilms tumor (WT) 1-specific cytotoxic T cells were observed in seven out of eight cancer patients who received a combination of WT1-peptide-pulsed DC-based

vaccine and S-1 or S-1 plus gemcitabine after surgery[59]. Of note, the procedure for developing DC vaccines is highly laborious and time-consuming and requires autologous cell preparations that do not meet economic requirement of precision therapy. Cell-based vaccines also include autologous and allogeneic tumor cell-derived vaccines[57,59]. Even though autologous tumor cell-based vaccines are particularly suitable for personalized therapy, autologous tumor cells may be insufficient, as only 15%–20% of pancreatic cancer patients are eligible for surgery[3,57]. Therefore, allogeneic tumor cell-based vaccines, including GVAX [allogeneic granulocyte-macrophage colony-stimulating factor (GM-CSF)-secreting

Table 1 Completed clinical trials of pancreatic cancer vaccines

Vaccine type	NCT number	Immunogen	Additional treatment	Phase	Enrollment	Endpoint
Cell-based vaccine	NCT00004604	CEA RNA-pulsed autologous DC	No	Phase I	Not provided	2002
	NCT00002773	Allogeneic pancreatic cancer cell	Cyclophosphamide, GM-CSF	Phase II	Not provided	2004
	NCT00084383	GVAX	5-fluorouracil, radiotherapy	Phase II	60	2006
	NCT00255827	Allogeneic tumor cell expressing α -1,3 galactosyltransferase	No	Phase I/II	7	2007
	NCT00128622	Autologous DC-infected with fowlpox-CEA-6D-TRICOM	Denileukin diftitox	Phase I	Not provided	2007
	NCT00027534	Autologous DC-infected with fowlpox-CEA-6D-TRICOM	Autologous DC mixed with CMV pp65 and tetanus toxoid	Phase I	Not provided	2007
	NCT00547144	Autologous DC	Gemcitabine, stereotactic radiosurgery	Phase I	2	2008
	NCT00002475	Allogeneic or autologous tumor cell	Cyclophosphamide, GM-CSF	Phase II	Not provided	2009
	NCT00305760	GVAX	Cetuximab, cyclophosphamide	Phase II	60	2009
	NCT00161187	Allogeneic lymphocyte	No	Phase I	Not provided	2011
	NCT01410968	Peptide-pulsed DC	Poly-ICLC	Phase I	12	2016
	NCT02151448	Autologous α DC1-loaded with autologous tumor material	Celecoxib, IFN- α , rintatolimod	Phase I/II	64	2019
	NCT00727441	GVAX	Surgery, cyclophosphamide	Phase II	87	2019
	NCT01896869	GVAX	FOLFIRINOX, ipilimumab	Phase II	83	2019
	Peptide-based vaccine	NCT00006387	RAS	Immunological adjuvant QS21	Phase I	Not provided
NCT00008099		MUC1	SB AS-2	Phase I	25	2004
NCT00019006		RAS	Detox-B adjuvant	Phase I	Not provided	Not provided
NCT00019331		RAS	IL-2, GM-CSF	Phase II	Not provided	2007
NCT00648102		HCG- β	No	Phase I	Not provided	2009
NCT00622622		VEGFR2	Gemcitabine	Phase I	21	2009
NCT00709462		HCG- β	No	Phase I	Not provided	2010
NCT00529984		CEA	No	Phase I/II	Not provided	2010
NCT00425360		Telomerase	Gemcitabine, capecitabine, GM-CSF	Phase III	Estimated 1110	2013
NCT00655785		VEGFR1, VEGFR2	Gemcitabine	Phase I/II	17	2013
NCT01342224	Telomerase	GM-CSF, gemcitabine	Phase I	11	2018	
Microorganism-based vaccine	NCT00003125	ALVAC-CEA, vaccinia-CEA	IL-2, GM-CSF	Phase II	Not provided	2004
	NCT00028496	Fowlpox-CEA(6D)-TRICOM	GM-CSF	Phase I	Not provided	2005
	NCT01191684	MVAp53	No	Phase I	Not provided	2013
	NCT00569387	Algenpantucel-L	Surgery, gemcitabine and 5-fluorouracil	Phase II	73	2014
	NCT00300950	Yeast expressing four different mutated RAS protein	Gemcitabine	Phase II	176	2015
	NCT02338752	DPT, typhoid, staphylococcus aureus, paratyphoid A and B	Surgery, chemotherapy	Phase I/II	20	2015
	NCT03127098	Adenovirus [E1-, E2b-]-CEA(6D)	IL-15	Phase I/II	Not provided	2017
Protein-based vaccine	NCT00003025	HSPPC-96	No	Phase I	16	2002
DNA-based vaccine	NCT01486329	VEGFR-2 DNA	No	Phase I	72	2014

All clinical trial data were collected from ClinicalTrials.gov (<https://clinicaltrials.gov/ct2/home>). CEA. Carcinoembryonic antigen; DC. Dendritic cell; GM-CSF. Granulocyte-macrophage colony-stimulating factor; GVAX. GM-CSF gene-transfected allogeneic pancreatic cancer cell; CMV pp65. Cytomegalovirus pp65; α DC1. α -type-1 polarized dendritic cell; RAS. Ras GTPase-activating protein; MUC1. Mucin 1; VEGFR. Vascular endothelial growth factor receptor; HCG- β . Human chorionic gonadotropin beta; SB AS-2. An immunologic adjuvant system consisting of an oil-in-water emulsion containing two immunostimulants: monophosphoryl Lipid A and a saponin derivative QS-21; MVAp53. Modified vaccinia virus ankara vaccine expressing p53; DPT. Diphtheria, pertussis, tetanus; HSPPC-96. Heat shock protein-peptide complex-96

Table 2 Ongoing clinical trials of pancreatic cancer vaccines

Vaccine type	NCT number	Immunogen	Additional treatment	Phase	Estimated enrollment	Status	Start point
Cell-based vaccines	NCT00389610	GVAX	No	Phase II	56	Active, not recruiting	2006
	NCT01088789	GVAX	Cyclophosphamide	Phase II	72	Recruiting	2010
	NCT01595321	GVAX	SBRT, FOLFIRINOX, cyclophosphamide	Not applicable	19	Active, not recruiting	2012
	NCT02451982	GVAX	Cyclophosphamide	Phase II	76	Recruiting	2016
	NCT02648282	GVAX	Cyclophosphamide, pembrolizumab, SBRT	Phase II	58	Active, not recruiting	2016
	NCT03190265	GVAX	Cyclophosphamide, nivolumab, CRS-207, ipilimumab	Phase II	63	Active, not recruiting	2017
	NCT03161379	GVAX	SBRT, nivolumab, cyclophosphamide	Phase II	30	Active, not recruiting	2018
	NCT03592888	Autologous DC pulsed with mutant KRAS peptides	No	Phase I	12	Recruiting	2018
	NCT03006302	GVAX	Epacadostat, pembrolizumab, CRS-207, cyclophosphamide	Phase II	40	Active, not recruiting	2018
	NCT03153410	GVAX	Cyclophosphamide, pembrolizumab, IMC-CS4	Phase I	12	Active, not recruiting	2018
	NCT03767582	GVAX	SBRT, nivolumab, CCR2/CCR5 dual antagonist	Phase I/II	30	Recruiting	2019
	NCT04157127	Autologous DC loaded with tumor lysate plus mRNA	No	Phase I	43	Recruiting	2020
	NCT04627246	Autologous DC loaded with personalized peptides	Nivolumab, chemotherapy	Phase I	12	Recruiting	2020
	Peptide-based vaccines	NCT03558945	Personalized neoantigen	Poly-ICLC	Phase I	60	Recruiting
NCT04161755		Personalized neoantigen	Atezolizumab, surgery, FOLFIRINOX	Phase I	29	Active, not recruiting	2019
NCT04117087		KRAS	Nivolumab, ipilimumab	Phase I	30	Recruiting	2020
NCT03956056		Personalized neoantigen and mesothelin	Poly-ICLC	Phase I	12	Active, not recruiting	2020
NCT04810910		Personalized neoantigen	Surgery, chemotherapy	Phase I	20	Recruiting	2021
NCT05111353		Neoantigen synthetic long peptide	Poly-ICLC	Phase I	30	Not yet recruiting	2022
NCT05013216		KRAS	Poly-ICLC	Phase I	25	Recruiting	2022
Microorganism-based vaccines	NCT00669734	Vaccinia, fowlpox	GM-CSF	Phase I	18	Active, not recruiting	2010
	NCT03136406	Recombinant saccharomyces cerevisiae yeast expressing mutant Ras	Cyclophosphamide, oxaliplatin, GI-4000, capecitabine, 5-fluorouracil, leucovorin, nab-paclitaxel, aNK, bevacizumab, avelumab, ALT-803, ETBX-011	Phase I/II	3	Active, not recruiting	2017
	NCT05116917	Influenza virus	Nivolumab, ipilimumab, SBRT	Phase II	30	Recruiting	2021
DNA-based vaccines	NCT03122106	Personalized neoantigens and mesothelin DNA	No	Phase I	15	Active, not recruiting	2018

All clinical trial data were collected from ClinicalTrials.gov (<https://clinicaltrials.gov/ct2/home>). GM-CSF. Granulocyte-macrophage colony-stimulating factor; GVAX. GM-CSF gene-transfected allogeneic pancreatic cancer cell; SBRT. Stereotactic body radiation therapy; CRS-207. *Listeria monocytogenes*-expressing mesothelin; DC. Dendritic cell; KRAS. GTPase KRas; CCR. C-C chemokine receptor; aNK. NK-92 cells

pancreatic cancer vaccine] and Algenpantucel-L (hyperacute-pancreatic cancer vaccine), are alternatives for pancreatic cancer treatment[57,60,61]. However, this approach does not consider the extensive heterogeneity of pancreatic cancer and thus is not suitable for personalized therapy. Compared with cyclophosphamide, a phase II trial revealed that GVAX did not improve the survival of patients with metastatic PDAC[60]. In one multi-institutional phase II clinical trial, the 12-month overall survival and disease-free survival rate of pancreatic cancer patients after treatment with Algenpantucel-L combined with standard adjuvant chemoradiotherapy reached 86% and 62%, respectively[60,62].

Microorganism-based pancreatic cancer vaccines

Microorganism-based pancreatic cancer vaccines are classified into bacteria, viruses, and recombinant yeast-based forms[63,64]. These vaccines represent a coexpressing strategy of tumor antigens and costimulatory molecules. The human adenovirus 40-based mesothelin vaccine inhibited the growth and metastasis of pancreatic cancer in mice[65]. An open-label phase I study of advanced pancreatic cancer showed that recombinant prime-boost poxviruses (targeting MUC1 and carcinoembryonic antigen prolonged the overall survival of patients with anti-MUC1 and/or carcinoembryonic antigen-specific immune responses[66]. In contrast, a phase II clinical trial revealed that compared with chemotherapy, vaccination with live-attenuated *Listeria monocytogenes* expressing mesothelin had no significant overall survival benefits for metastatic PDAC patients[67]. Notably, microorganism-based vaccines require complicated engineering system and elaborate fabrication, undermining their inconvenient application for personalized treatment.

Exosome-based pancreatic cancer vaccines

Tumor-derived exosomes (TEXs) are nanosized lipid bilayer encapsulating vesicles that shuttle bioactive information to the tumor microenvironment, promoting tumor progression[68,69]. TEXs contain various tumor antigens and feature discrete sets of specific proteins that promote DC-binding and uptake of exosomes. In pancreatic cancer mouse models, DCs loaded with TEXs vaccine activated CD4⁺ T cells and significantly prolonged the survival of mice compared to cytotoxic drugs[70]. Notably, TEXs also include proteins and nucleic acids which have strong capability to boost the body's immunity, and thus may cause auto-immune diseases by disrupting the immune homeostasis after vaccination, posing a challenge to safety of precision therapy[71-74].

Protein-based pancreatic cancer vaccines

Proteins for cancer vaccination are not only immunogenic but also can carry additional antigenic peptide. Proteins vaccines

based on heat shock proteins, especially heat shock protein-peptide complex-96 (HSPPC-96), are currently under several clinical trials to investigate their therapeutic potential against different cancers[75-80]. A phase I pilot study revealed that 30% (3/10) of patients with resected pancreatic cancer survived for more than 5 years after vaccination with the HSPPC-96 vaccine[75]. Notably, given that HSPPC-96 must be extracted from tumor tissues of each patient, its use largely depends on the resectability of the tumors[80].

Peptide-based pancreatic cancer vaccines

Peptide-based vaccines are developed based on antigenic epitopes, the minimal immunogenic regions of antigens[81]. KRAS-targeting peptide was the first peptide-based vaccine to undergo clinical trials[82]. In a phase I/II study, GM-CSF combined with KRAS-targeting peptide vaccine-induced specific immune response in 25 of 43(58%) patients, and the survival period was also significantly longer for responders than non-responders[83]. Another commonly tested peptide-based vaccine, the telomerase-targeting vaccine (GV1001), was well tolerated and improved patient survival in a phase I/II clinical trial[84]. However, two phase III clinical trials revealed that compared with mono-gemcitabine, a combination of GV1001 with gemcitabine did not significantly improve the overall survival of patients with advanced pancreatic cancer[85]. Aside from KRAS and telomerase-targeted peptides, clinical trials have revealed that the efficacy of survivin, gastrin, vascular endothelial growth factor receptor (VEGFR)-1, VEGFR-2, WT1, and kinesin family member 20A-targeted vaccines is unsatisfactory[86-90]. Notably, the tumor peptide vaccine is major histocompatibility complex (MHC)-restricted and only activates monoclonal T cells, which may reduce the strength of anti-tumor immune response and thus do not satisfy the need of efficiency underlying precision therapy[81].

DNA-based pancreatic cancer vaccines

DNA-based vaccines serve as templates encoding antigens in transfected cells. Enolase 1 (ENO1), MUC1, survivin, and VEGFR-2-targeting DNA vaccines are examples of the DNA-based pancreatic cancer vaccines explored so far[91-95]. Preclinically, the ENO1 DNA vaccine efficiently induced the infiltration of effector T cells, antibody formation, and tumor cytotoxicity in genetically engineered mice with pancreatic cancer[92]. Moreover, combined with chemotherapy, the ENO1 DNA vaccine induced CD4⁺ T cell-mediated antitumor activity and strongly impaired cancer progression in mice[92]. MUC1-targeted DNA vaccine induced strong and specific cytotoxic T lymphocyte response and showed both therapeutic and prophylactic effects in mice[91]. The survivin DNA vaccine induced specific antitumor immunity

and prolonged the survival period of mice[94]. Also, VXM01, an oral DNA vaccine targeting VEGFR-2, is under phase I trial for stage IV pancreatic cancer treatment[95]. Notably, DNA vaccines increase the risk of host genomic alteration, the coded antigens are expressed over a long-time, and the production of anti-DNA autoantibodies may limit their application[96,97]. Obviously, the safety is a major concern for application of DNA vaccine in personalized pancreatic cancer treatment.

In summary, these conventional vaccines show a measure of progress in pancreatic cancer therapy. However, given the major concerns, including the safety and complexity in preparation, they are not the best options for vaccines-based personalized precise treatment of pancreatic cancer. Therefore, it is essential to select a novel kind of vaccines that meet the needs of individual pancreatic cancer patients.

Superiority of mRNA vaccine for cancer precision treatment

mRNA vaccines are emerging as potent candidates for cancer precision treatment because of their unique advantages over the above-mentioned vaccine formats. In addition to overcoming tumor heterogeneity by encoding personalized protein according to the genetic expression profile of tumor, mRNA vaccine meets the requirements of precision therapy highlighting precise targeting, high efficiency, safety, and economic cost.

Generation of natural protein products

As fore-mentioned, precise targeting is key for personalized therapy. Consistent with this point, mRNA functions as a template for protein translation, and utilizes the machinery in host cells for vaccine production. This characteristic allows for post-translational modification of the protein products, including proper folding for effective functioning[44,98]. Also, this approach allows for the production of correctly folded and assembled multimeric proteins that cannot be generated in bioreactors; this method allows for the produced transmembrane and intracellular proteins to be translocated to the appropriate specific cellular sites. Therefore, mRNA vaccine generates protein products with endogenous characteristics, ensuring the precision of targeting.

Induction of both innate and adaptive immunities

Efficiency is another keypoint in precision therapy. Meeting the requirement, mRNA vaccine can induce both innate and adaptive immunities to exert efficient anti-tumor effects. Innate immunity forms the first line of defense against non-self antigens[97]. APCs, especially DCs, engulf foreign mRNA *via* pattern recognition receptors (PRRs), activating a series of proinflammation-related signaling pathways that promotes

the function of innate immunity[97,99]. For example, PRR Toll-like receptor (TLR)-3 recognizes and binds double strand RNA, regulating the secretion of cytokines and chemokines as well as the activation of the type I interferon (IFN) pathway[100]. In addition, PRR TLR-7 and TLR-8 bind single-strand RNA, activating nitric oxide synthase and the production of type I IFN[101-103]. The secretion of type I IFN is essential for the formation of an immune-stimulatory environment, wherein T cells differentiate into cytotoxic types that can eliminate tumors. Apart from innate immunity, mRNA vaccines further stimulate adaptive immunity. The protein encoded by non-self mRNA can be degraded into peptides, which are routed into the endoplasmic reticulum, loaded onto MHC-I, shuttled to the cell surface, and ultimately presented to and activate CD8⁺ T cells[98,104,105]. Meanwhile, the antigens can be transported from Golgi to endosomes and enter the MHC-II presentation pathway, where they activate CD4⁺ T cells[106]. Actually, the antigens can also be secreted and reinternalized and presented *via* MHC-II to activate CD4⁺ T cells or cross-presented *via* MHC-I to activate CD8⁺ T cells[104,106]. Moreover, mRNA vaccine can upregulate the expression of costimulatory molecules (e.g., CD40 and CD86) on APCs (e.g., DCs), enhancing the antigen presentation and T cell activation[107]. Furthermore, activated APCs (e.g., macrophage and DC) present antigens to activate B cells, triggering an antibody response[108,109]. Multiple preclinical and clinical trials have shown that mRNA vaccines induce antitumor immune responses and tumor rejection. Melanoma mouse models have revealed that mRNA-lipoplexes encoding mutant or viral neo-antigens or endogenous self-antigens trigger IFN- α release by macrophages and plasmacytoid DCs, induce strong effector and memory T-cell responses, and mediate the rejection of progressive tumors[107]. A personalized mRNA vaccine induces T cell infiltration and specific killing of melanoma[43]. Additionally, the intravenously administered liposomal RNA vaccine BNT111 mediates a durable objective response and induces strong anti-melanoma CD4⁺ and CD8⁺ T cell immunity after pretreatment with an immune checkpoint inhibitor[110]. In summary, mRNA vaccines accord with the concerns about efficiency in precision therapy, inducing both innate immunity and adaptive immunity to exert potent anti-tumor effects.

High safety in practice

Safety is also important for precision therapy. In line with this, mRNA production does not involve toxic chemicals and the risk of contamination with the adventitious virus packaged in cell cultures. Therefore, an mRNA approach averts common threats associated with other vaccine platforms (e.g., viral

vectors, inactivated viruses, live viruses, and subunit protein vaccines). In addition, the rapid manufacturability of mRNA decreases opportunities for the introduction of contaminating microorganisms. In this context, it is also important to note that mRNA cannot integrate into the host genome, ruling out oncogenic potentials. Finally, mRNA can be rapidly degraded by RNA enzymes and is characterized by its adjustable half-life, which defines the controllable expression of mRNA-encoded proteins[111-113]. The first clinical study of mRNA vaccine was conducted in 2008 in melanoma patients[114]. Vaccination with naked mRNA is safe and well tolerated and does not induce World Health Organization grade III or IV adverse events. Numerous clinical trials have supported the high safety of mRNA vaccines[115-117]. For example, direct injection of protamine-protected mRNA into patients with metastatic melanoma predominantly caused local inflammatory skin reactions or fatigue, which could be easily lessened by symptomatic therapy. Therefore, mRNA vaccines are relatively safe, which is consistent with the principle of safety underlying precision therapy.

Convenience and low cost of preparation

Economic principle is last, but perhaps most important, in precision therapy. Exactly, preparation of mRNA vaccine is convenient and low cost. mRNA can be produced *in vitro* using a DNA template, ribonucleotide triphosphates, and recombinant enzymes[118,119]. In this process, a plasmid DNA containing a DNA-dependent RNA polymerase promoter (e.g., T3, T7, or SP6) is first generated. It is then linearized to provide a template for mRNA synthesis using DNA-dependent RNA polymerase before degradation by DNase. A 5'cap and a 3'poly-A tail are added during the transcription step to facilitate efficient translation *in vivo*. Finally, free nucleotides, enzymes, truncated RNA fragments, and residual DNA are removed to obtain pure mRNA. This simple process ensures rapid mRNA production in a relatively less complex system and thus can be standardized to produce almost any encoded protein immunogen, rendering it highly suitable for constructing personalized vaccines for cancer treatment. Moreover, all reaction components and enzymes required for mRNA production are commercially available. The entire process of mRNA vaccine production takes about ten days, significantly shorter than other formats[53]. The rapid production of mRNA vaccine is a tremendous advantage for personalized therapy, meaning that treatment can be available within a short time after diagnosis. From an industrial perspective, the large-scale production of mRNA vaccines is lowcost. DNA templates are used during the transcription cycle and by scaling the *in vitro* transcription reaction. A very

small (about 1 μ g) DNA template can produce very large amounts (hundreds μ g) of capped mRNA, and the product is more dependent on the transcription volume and time than on the amount of DNA[120]. In addition, the required mRNA vaccine dose is generally lower than DNA vaccines (50–100 μ g for an mRNA vaccine and 1–5 mg for a DNA vaccine). Actually, an only 10 g of mRNA can generate about 100,000 vaccine doses. Together, the production of mRNA vaccines is convenient and low-cost, which is in agreement with the economic principle of precision therapy.

Development strategy of personalized mRNA vaccines for pancreatic cancer

With precise targeting, efficiency, safety, and economic cost, mRNA vaccines offer promise for providing personalized pancreatic cancer treatment. Accumulating evidence suggests that the pipeline for developing personalized pancreatic cancer mRNA vaccines should be divided into three critical modules, including identifying tumor antigens, constructing mRNA vaccines, and distinguishing immune subtypes (Fig. 2)[29].

Identification of pancreatic cancer antigens

Antigen selection is the first step in developing a vaccine. An ideal vaccine candidate should possess the following characteristics: unique to tumor cells, involved in tumorigenesis and progression, non-tolerated by the immune system, and stimulatory to the antitumor immunity[121-123]. Current immunogenic targets for cancer treatment include tumor-associated antigens (TAAs) and tumor-specific antigens (TSAs)[124]. TAAs are the most commonly targeted antigens typically expressed on normal cells but aberrantly on tumor cells. This underscores their potential as universal therapeutic targets, although they are self-antigens and thus may be immunologically tolerated and weakened vaccine potency. Unlike TAAs, TSAs are exclusively expressed in tumor cells with strong immunogenicity and a high degree of individuality and epitope diversity and thus are ideal targets for personalized vaccines. Engineering a personalized anti-pancreatic cancer mRNA vaccine begins with identifying tumor-specific non-synonymous mutations by comparing next-generation sequencing data of tumors and paired normal tissues[125]. Computational neoantigen prediction pipelines are then applied to verify the expression and predict the binding affinity of peptides generated from mutated genes onto MHC alleles. High transcript expression is related to enhanced T cell response and can compensate for the low MHC-binding affinity of mutations[126]. Furthermore, a single MHC-I-bound TSA is not sufficient, and additional MHC-II-bound TSAs are needed for effective antitumor immunity[127].




Fig. 2 Streamlined development of personalized mRNA vaccines for pancreatic cancer.

Novel tumor antigens are identified as potent targets for the preparation of promising pancreatic cancer mRNA vaccines. Immune subtypes are identified as vital criteria for selecting applicable pancreatic cancer patients for mRNA vaccine treatment. Partial elements of this figure are adopted from Huang et al.[29] with appropriate modification. ORF. Open reading frame; DC. Dendritic cell; APCs. Antigen-presenting cells; OS. Overall survival; RFS. Relapse-free survival

NetMHCpan and MHCflurry are tools trained for predicting the binding affinity between ligands and MHC[128-130]. Notably, for predicting immunogenicity, the stability of the

neopeptide-MHC complex is more important than the binding affinity[131]. NetMHCstabpan, a tool for stability prediction, performs well in identifying immunogenic mutations[132]. In

addition to the surface presentation, the interaction between peptide-MHC complex and T-cell receptor is necessary to induce an immune response, and predicting this interaction is based on amino acid side chains of the T-cell receptor facing the MHC-bound peptide[133]. Recently, a perspective pipeline for identifying tumor antigens by screening for the over-expressed and mutated genes and prognosis and APC-associated candidates has been established[29]. Notably, although the above-mentioned characteristics are based upon a sound rationale, a specific approach to weigh each of them has not been set up, and therefore, optimal candidates for mRNA vaccine development cannot be selected. Nevertheless, tumor antigen prediction is rapidly evolving thanks to the recent progress in computational biology. Accordingly, an accurate and sensitive approach for identifying potent candidates in individual pancreatic cancer will eventually be established for developing its personalized mRNA vaccines.


Construction of mRNA vaccines against pancreatic cancer

Several critical issues, including delivery, stability, translation, and immunogenicity, must be addressed before the practical application of the mRNA-based cancer vaccine[97,98,134,135]. Because of its size, degradability, and charge, naked mRNA cannot efficiently pass through the cell membrane and enter the cytoplasm, except for immature DCs that can efficiently uptake mRNA *via* the macro-pinocytosis pathway[134,136]. For more effective delivery of mRNA into APCs, mRNA formulations (e.g., liposomes, polyplexes, polysomes, and lipoplexes) and administration routes must be appropriately selected and optimized. After successful mRNA delivery, the half-life of mRNA transcribed *in vivo* must be appropriately regulated, given that several factors influence the pharmacodynamic and pharmacokinetic properties of mRNA-based therapeutics. There is a need to improve mRNA structures, including optimization of poly (A), 5'cap, poly-A tail, untranslated regions, and protein encoding open reading frames, to enhance the stability of mRNAs[115,137,138]. In addition to delivery and stability, immunogenicity must also be considered. Accumulating evidence suggests that there is a negative feedback loop between mRNA and it-induced immune response. For instance, exogenous RNA stimulates the production of type I IFN by stimulating innate immunity[98], while excessive production of type I IFN inhibits translation and promotes the degradation of both ribosomal RNA and cellular mRNA[98,134,139,140]. The addition of poly-A tails, optimization of sequences, and posttranscriptional purification can decrease the level of innate immunity without altering the translation of mRNA[112,141-145]. Furthermore, increasing the immunostimulatory properties of mRNA using adjuvants

promotes the potency of cancer mRNA vaccines. TriMix (mRNA encoding CD70, CD40L, and TLR4) enhances the immunogenicity of unmodified naked mRNA and improves the cytotoxicity of T lymphocyte and DC maturation[98]. Together, advances in optimization strategies for mRNA vaccine construction largely improve the efficacy of these vaccines for pancreatic cancer treatment.

Distinction of immune subtypes in pancreatic cancer

Pancreatic cancer is usually characterized by a complex immunosuppressive microenvironment, low mutational burden, and poor T cell infiltration[11,146,147]. Although an mRNA vaccine can activate and promote infiltration of T cells into the tumor, the entry of these cells could still be largely interfered with by the desmoplastic stroma of pancreatic cancer cells[4,148]. Moreover, numerous immunosuppressive cells (e.g., myeloid cells, regulatory T cells, and M2 macrophages), signaling pathways (e.g., transforming growth factor beta signaling pathway, IL-10 signaling pathway, and VEGF signaling pathway), and molecules (e.g., PD-L1, T cell immunoglobulin mucin 3, T cell immunoreceptor with Ig and ITIM domains, lymphocyte activating 3, V-type immunoglobulin domain-containing suppressor of T-cell activation, and CD73) lead to multiple immunosuppression on anti-tumor immune response in the pancreatic cancer microenvironment[11,146]. Hence, mRNA vaccines in combination with other therapies (rather than a single vaccine) and biomarkers for predicting therapeutic response of combination strategies are strongly needed for pancreatic cancer treatment. To date, diverse pancreatic cancer subtypes, defined based on different parameters, approaches, and perspectives, have been identified (Fig. 3)[28,29,31,149-159]. Multiple immunological factors, including immune-related gene expression profile and immune cell composition, are used for grouping immune subtypes of pancreatic cancer. Immune subtypes indicate the immunological status in pancreatic tumors and their microenvironment and thus are accurate biomarkers for selecting a suitable combined therapy[29,149,150]. For instance, immunologically "cold" pancreatic tumors generally show low immunogenicity and/or high reactive stroma, whereas immunogenic chemotherapy and stromal modulation may promote the effectiveness of an mRNA vaccine by improving tumor immunogenicity and T cell infiltration. Additionally, an mRNA vaccine combined with immune checkpoint blockade may improve T cell infiltration and function in immunologically "hot" tumors. Overall, combination therapy may enhance the efficacy of an mRNA vaccine for pancreatic cancer treatment under the guidance of immune subtypes, biomarkers for matching patients and therapeutics.




Concluding remarks and outlook

The mRNA vaccine is a novel and promising vehicle for developing personalized vaccines against pancreatic cancer. Identification of potent tumor antigens, optimization of immunostimulatory vaccine construction, and distinction of immune subtypes are prerequisites for the personalization of potentially effective pancreatic cancer mRNA vaccines.

At present, a first-in-human phase I study on the tolerability and safety of the mRNA-based personalized neoantigen vaccine (autogene cevumeran, also known as BNT122, RO7198457) in combination with chemotherapy and PD-L1 blockade for resected PDAC is currently underway. The preliminary findings in this trial were released for the first time on June 5, 2022 by BioNTech Company (<https://investors.biontech.de/>). Sixteen patients who underwent surgery and received PD-L1 inhibition were vaccinated with autogene cevumeran, and all well tolerated the treatment. Only one developed a vaccine-related grade three fever and hypertension, and no other grade three or higher adverse events were observed. In addition, half were detected de-novo neoantigen-specific T cell responses, and had a significantly longer recurrence-free survival (median not determined, but is more than 18 months) compared with those without vaccine-induced immune responses (13.4 months). Therefore, personalized mRNA vaccine is a promising strategy for pancreatic cancer treatment. Notably, only about 15%–20% of pancreatic cancer patients present with localized disease that can be resected through the standard procedure[3]. In contrast, the majority of pancreatic cancers are diagnosed at the locally advanced or metastatic level and/or are poorly differentiated and ineligible for surgery, rendering the acquirement of these tumors

largely dependent on biopsy. However, the inter- and intra-tumoral heterogeneity of pancreatic cancer limits the actual value of biopsy-derived samples, at least not sufficient for the personalized design and construction of mRNA vaccines as well as the distinction of patients for suitable combination therapy. Neoadjuvant therapies can be used to reduce tumor staging and eliminate micro-metastases, increasing the chance of successful surgery[160-162]. For instance, after treatment with the neoadjuvant FOLFIRINOX, 76 of 125(60.8%) patients with unresectable pancreatic cancer qualified for tumor resection[160]. In a separate study, 141 patients with unresectable (51.1%) or borderline-resectable (48.9%) non-metastatic cancers were recruited; of these, 78% qualified and underwent surgery after FOLFIRINOX therapy[161]. Therefore, neoadjuvant therapies may facilitate the acquisition of relatively sufficient tumor samples for identifying individualized tumor antigens and immune subtypes for developing personalized mRNA vaccines.

Nevertheless, treating pancreatic cancer with mRNA vaccines remains challenging. As mentioned above, pancreatic cancer is highly heterogeneous. Due to the complexity of the pancreatic tumor components, including both therapeutically sensitive cancer cells and genetically resistant cancer cells or epigenetically plastic persister cancer cells, the therapeutic stress may cause tumor evolution and consequent treatment failure (Fig. 4). Alternatively, the development of prophylactic mRNA vaccines may be another considerable strategy against this intractable disease. To date, prophylactic mRNA vaccination is mainly applied for preventing infection of viruses, since they are ectogenic and possess simple construction and antigens for vaccine development can easily



be identified. In contrast, the development of prophylactic vaccines against pancreatic cancer is still in infancy, partially but not totally, due to the complexity of pancreatic cancer onset and the difficulty of assessing the effectiveness of vaccination. Research shows that PDAC arises from noninvasive precancerous lesions, microscopic pancreatic intraepithelial neoplasia (PanIN), and macroscopic intraductal papillary mucinous neoplasms (IPMNs)[163]. In processing PanIN-PDAC transformation, oncogenic mutation in KRAS gene has been detected in >90% of the low-grade-PanINs[163]. In IPMN-PDAC progression, oncogenic mutations in KRAS and GNAS genes have been observed in 50%–80% and 40%–70% of IPMN, respectively[164]. KRAS and GNAS are therefore considered potential targets for preventing PDAC development. A KRAS-targeting peptide vaccine for preventing pancreatic cancer in high-risk individuals is currently under clinical test (NCT05013216), but its prophylactic efficiency is still unknown. Notably, apart from KRAS mutation, multiple genetic and epigenomic mechanisms jointly contribute to pancreatic cancer initiation. For instance, IL-33 is identified as a key factor that induces epigenetic reprogramming-mediated pancreatic oncogenesis[165,166]. The question that whether cotargeting KRAS and IL-33 increases the probability of pancreatic cancer prevention arises and remains to be validated.

Abbreviations

APC: Antigen-presenting cell; COVID-19: Coronavirus disease 2019;

DC: Dendritic cell; ENO1: Enolase 1; GM-CSF: Granulocyte-macrophage colony-stimulating factor; HSPPC-96: Heat shock protein-peptide complex-96; IFN: Type I interferon; IPMNs: Intraductal papillary mucinous neoplasms; MHC: Major histocompatibility complex; mRNA: Messenger RNA; MUC1: Mucin 1; PanIN: Pancreatic intraepithelial neoplasia; PDAC: Pancreatic ductal adenocarcinoma; PD-1: Programmed cell death 1; PD-L1: Programmed cell death ligand 1; PPRs: Pattern recognition receptors; SARS-CoV-2: Severe acute respiratory syndrome coronavirus 2; TAAs: Tumor-associated antigens; TEXs: Tumor-derived exosomes; TLR: Toll like receptor; TSAs: Tumor-specific antigens; VEGFR: Vascular endothelial growth factor receptor; WT1: Wilms tumor 1.

Acknowledgements

The author Xing Huang would like to convey his sincerest and deepest gratitude to Prof. Guido Kroemer (Centre de Recherche des Cordeliers) for the technological training and ideological inspiration in his laboratory.

Authors' contributions

XH, GZ, and TBL conceived the review. XH and GZ drafted and revised the manuscript. TYT and XG helped for proof-reading. All authors discussed and approved the final paper.

Funding

This work was supported by the National Natural Science Foundation of China (31970696, 81502975, 82188102, and 81830089), Zhejiang Provincial Natural Science Foundation for Distinguished Young Scholar (LR22H160010), National Key Research and Development Program of China (2019YFC1316000), Zhejiang Provincial Key Research and Development Program (2019C03019), and Zhejiang Provincial College Student Science and Technology Innovation Activity Plan-College Student Innovation and Entrepreneurship Incubation Program (Young

Talent Program) (2022R40122).

Availability of data and materials

Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Zhejiang Provincial Key Laboratory of Pancreatic Disease, the First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310009, China. ²Department of Hepatobiliary and Pancreatic Surgery, the First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China. ³Zhejiang Clinical Research Center of Hepatobiliary and Pancreatic Diseases, Hangzhou 310003, China. ⁴The Innovation Center for the Study of Pancreatic Diseases of Zhejiang Province, Hangzhou 310009, China. ⁵Cancer Center, Zhejiang University, Hangzhou 310058, China.

References

1. Klein AP. Pancreatic cancer epidemiology: understanding the role of lifestyle and inherited risk factors. *Nat Rev Gastroenterol Hepatol.* 2021;18(7):493–502.
2. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2021;71(3):209–49.
3. Strobel O, Neoptolemos J, Jäger D, Büchler MW. Optimizing the outcomes of pancreatic cancer surgery. *Nat Rev Clin Oncol.* 2019;16(1):11–26.
4. Hessmann E, Buchholz SM, Demir IE, Singh SK, Gress TM, Ellenrieder V, et al. Microenvironmental determinants of pancreatic cancer. *Physiol Rev.* 2020;100(4):1707–51.
5. Conroy T, Hammel P, Hebbbar M, Ben Abdelghani M, Wei AC, Raoul JL, et al. FOLFIRINOX or gemcitabine as adjuvant therapy for pancreatic cancer. *N Engl J Med.* 2018;379(25):2395–406.
6. Von Hoff DD, Ervin T, Arena FP, Chiorean EG, Infante J, Moore M, et al. Increased survival in pancreatic cancer with nab-paclitaxel plus gemcitabine. *N Engl J Med.* 2013;369(18):1691–703.
7. Neoptolemos JP, Kleeff J, Michl P, Costello E, Greenhalf W, Palmer DH. Therapeutic developments in pancreatic cancer: current and future perspectives. *Nat Rev Gastroenterol Hepatol.* 2018;15(6):333–48.
8. Li E, Huang X, Zhang G, Liang T. Combinational blockade of MET and PD-L1 improves pancreatic cancer immunotherapeutic efficacy. *J Exp Clin Cancer Res.* 2021;40(1):279.
9. Zhang X, Huang X, Xu J, Li E, Lao M, Tang T, et al. NEK2 inhibition triggers anti-pancreatic cancer immunity by targeting PD-L1. *Nat Commun.* 2021;12(1):4536.
10. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2021. *Cancer J Clin.* 2021;71(1):7–33.
11. Bear AS, Vonderheide RH, O'Hara MH. Challenges and opportunities for pancreatic cancer immunotherapy. *Cancer Cell.* 2020;38(6):788–802.
12. Samuel N, Hudson TJ. The molecular and cellular heterogeneity of pancreatic ductal adenocarcinoma. *Nat Rev Gastroenterol Hepatol.* 2011;9(2):77–87.
13. Makohon-Moore AP, Zhang M, Reiter JG, Bozic I, Allen B, Kundu D, et al. Limited heterogeneity of known driver gene mutations among the metastases of individual patients with pancreatic cancer. *Nat Genet.* 2017;49(3):358–66.
14. Yachida S, White CM, Naito Y, Zhong Y, Brosnan JA, Macgregor-Das AM, et al. Clinical significance of the genetic landscape of pancreatic cancer and implications for identification of potential long-term survivors. *Clin Cancer Res.* 2012;18(22):6339–47.
15. Wang F, Xia X, Yang C, Shen J, Mai J, Kim HC, et al. SMAD4 gene mutation renders pancreatic cancer resistance to radiotherapy through promotion of autophagy. *Clin Cancer Res.* 2018;24(13):3176–85.
16. Devico Marciano N, Kroening G, Dayyani F, Zell JA, Lee FC, Cho M, et al. BRCA-mutated pancreatic cancer: from discovery to novel treatment paradigms. *Cancers.* 2022;14(10):2453.
17. Emelyanova M, Pudova E, Khomich D, Krasnov G, Popova A, Abramov I, et al. Platinum-based chemotherapy for pancreatic cancer: impact of mutations in the homologous recombination repair and Fanconi anemia genes. *Ther Adv Med Oncol.* 2022;14:17588359221083050.
18. Edwards SL, Brough R, Lord CJ, Natrajan R, Vatcheva R, Levine DA, et al. Resistance to therapy caused by intragenic deletion in BRCA2. *Nature.* 2008;451(7182):1111–5.
19. Chen A. PARP inhibitors: its role in treatment of cancer. *Chin J Cancer.* 2011;30(7):463–71.
20. Golan T, Hammel P, Reni M, Van Cutsem E, Macarulla T, Hall MJ, et al. Maintenance olaparib for germline BRCA-mutated metastatic pancreatic cancer. *N Engl J Med.* 2019;381(4):317–27.
21. Javle M, Shacham-Shmueli E, Xiao L, Varadhachary G, Halpern N, Fogelman D, et al. Olaparib monotherapy for previously treated pancreatic cancer with DNA damage repair genetic alterations other than germline BRCA variants: findings from 2 phase 2 nonrandomized clinical trials. *JAMA Oncol.* 2021;7(5):693–9.
22. Ribas A, Wolchok JD. Cancer immunotherapy using checkpoint blockade. *Science.* 2018;359(6382):1350–5.
23. Le DT, Uram JN, Wang H, Bartlett BR, Kemberling H, Eyring AD, et al. PD-1 blockade in tumors with mismatch-repair deficiency. *N Engl J Med.* 2015;372(26):2509–20.
24. Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SAJR, Behjati S, Biankin AV, et al. Signatures of mutational processes in human cancer. *Nature.* 2013;500(7463):415–21.
25. Le DT, Durham JN, Smith KN, Wang H, Bartlett BR, Aulakh LK, et al. Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science.* 2017;357(6349):409–13.
26. Stromnes IM, Hulbert A, Pierce RH, Greenberg PD, Hingorani SR. T-cell localization, activation, and clonal expansion in human pancreatic ductal adenocarcinoma. *Cancer Immunol Res.* 2017;5(11):978–91.
27. Huang X, Zhang G, Liang T. Subtyping for pancreatic cancer precision therapy. *Trends Pharmacol Sci.* 2022;43(6):482–94.
28. Moffitt RA, Marayati R, Flate EL, Volmar KE, Loeza SG, Hoadley KA, et al. Virtual microdissection identifies distinct tumor- and stromaspecific subtypes of pancreatic ductal adenocarcinoma.

- Nat Genet. 2015;47(10):1168–78.
29. Huang X, Tang T, Zhang G, Liang T. Identification of tumor antigens and immune subtypes of pancreatic adenocarcinoma for mRNA vaccine development. *Mol Cancer*. 2021;20(1):44.
 30. Peng J, Sun BF, Chen CY, Zhou JY, Chen YS, Chen H, et al. Singlecell RNA-seq highlights intra-tumoral heterogeneity and malignant progression in pancreatic ductal adenocarcinoma. *Cell Res*. 2019;29(9):725–38.
 31. Chan-Seng-Yue M, Kim JC, Wilson GW, Ng K, Figueroa EF, O’Kane GM, et al. Transcription phenotypes of pancreatic cancer are driven by genomic events during tumor evolution. *Nat Genet*. 2020;52(2):231–40.
 32. Huang X, Tang T, Zhang G, Liang T. Identification of tumor antigens and immune subtypes of cholangiocarcinoma for mRNA vaccine development. *Mol Cancer*. 2021;20(1):50.
 33. Negahdaripour M, Golkar N, Hajighahramani N, Kianpour S, Nezafat N, Ghasemi Y. Harnessing self-assembled peptide nanoparticles in epitope vaccine design. *Biotechnol Adv*. 2017;35(5):575–96.
 34. Brenner S, Jacob F, Meselson M. An unstable intermediate carrying information from genes to ribosomes for protein synthesis. *Nature*. 1961;190:576–81.
 35. Lockard RE, Lingrel JB. The synthesis of mouse hemoglobin chains in a rabbit reticulocyte cell-free system programmed with mouse reticulocyte 9S RNA. *Biochem Biophys Res Commun*. 1969;37(2):204–12.
 36. Wolff JA, Malone RW, Williams P, Chong W, Acsadi G, Jani A, et al. Direct gene transfer into mouse muscle *in vivo*. *Science*. 1990;247(4949 Pt 1):1465–8.
 37. Jirikowski GF, Sanna PP, Maciejewski-Lenoir D, Bloom FE. Reversal of diabetes insipidus in Brattleboro rats: intrahypothalamic injection of vasopressin mRNA. *Science*. 1992;255(5047):996–8.
 38. Martinon F, Krishnan S, Lenzen G, Magné R, Gomard E, Guillet JG, et al. Induction of virus-specific cytotoxic T lymphocytes *in vivo* by liposome-entrapped mRNA. *Eur J Immunol*. 1993;23(7):1719–22.
 39. Conry RM, LoBuglio AF, Loechel F, Moore SE, Sumerel LA, Barlow DL, et al. A carcinoembryonic antigen polynucleotide vaccine has *in vivo* antitumor activity. *Gene Ther*. 1995;2(1):59–65.
 40. Conry RM, LoBuglio AF, Loechel F, Moore SE, Sumerel LA, Barlow DL, et al. A carcinoembryonic antigen polynucleotide vaccine for human clinical use. *Cancer Gene Ther*. 1995;2(1):33–8.
 41. Hajj KA, Whitehead KA. Tools for translation: non-viral materials for therapeutic mRNA delivery. *Nat Rev Mater*. 2017;2:17056.
 42. Kim J, Eygeris Y, Gupta M, Sahay G. Self-assembled mRNA vaccines. *Adv Drug Deliv Rev*. 2021;170:83–112.
 43. Sahin U, Derhovannessian E, Miller M, Kloke BP, Simon P, Lower M, et al. Personalized RNA mutanome vaccines mobilize poly-specific therapeutic immunity against cancer. *Nature*. 2017;547(7662):222–6.
 44. Barbier AJ, Jiang AY, Zhang P, Wooster R, Anderson DG. The clinical progress of mRNA vaccines and immunotherapies. *Nat Biotechnol*. 2022;40(6):840–54.
 45. Shin MD, Shukla S, Chung YH, Beiss V, Chan SK, Ortega-Rivera OA, et al. COVID-19 vaccine development and a potential nanomaterial path forward. *Nat Nanotechnol*. 2020;15(8):646–55.
 46. Chung JY, Thone MN, Kwon YJ. COVID-19 vaccines: the status and perspectives in delivery points of view. *Adv Drug Deliv Rev*. 2021;170:1–25.
 47. Pascolo S. Vaccines against COVID-19: priority to mRNA-based formulations. *Cells*. 2021;10(10):2716.
 48. Dagan N, Barda N, Kepten E, Miron O, Perchik S, Katz MA, et al. BNT162b2 mRNA Covid-19 vaccine in a nationwide mass vaccination setting. *N Engl J Med*. 2021;384(15):1412–23.
 49. Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *N Engl J Med*. 2020;383(27):2603–15.
 50. Baden LR, El Sahly HM, Essink B, Kotloff K, Frey S, Novak R, et al. Efficacy and safety of the mRNA-1273 SARS-CoV-2 vaccine. *N Engl J Med*. 2021;384(5):403–16.
 51. Ali K, Berman G, Zhou H, Deng W, Faughnan V, Coronado-Voges M, et al. Evaluation of mRNA-1273 SARS-CoV-2 vaccine in adolescents. *N Engl J Med*. 2021;385(24):2241–51.
 52. Thomas SJ, Moreira ED Jr, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine through 6 months. *N Engl J Med*. 2021;385(19):1761–73.
 53. Fang E, Liu X, Li M, Zhang Z, Song L, Zhu B, et al. Advances in COVID-19 mRNA vaccine development. *Signal Transduct Target Ther*. 2022;7(1):94.
 54. Chemaitelly H, Yassine HM, Benslimane FM, Al Khatib HA, Tang P, Hasan MR, et al. mRNA-1273 COVID-19 vaccine effectiveness against the B.1.1.7 and B.1.351 variants and severe COVID-19 disease in Qatar. *Nat Med*. 2021;27(9):1614–21.
 55. Choi A, Koch M, Wu K, Chu L, Ma L, Hill A, et al. Safety and immunogenicity of SARS-CoV-2 variant mRNA vaccine boosters in healthy adults: an interim analysis. *Nat Med*. 2021;27(11):2025–31.
 56. Tang P, Hasan MR, Chemaitelly H, Yassine HM, Benslimane FM, Al Khatib HA, et al. BNT162b2 and mRNA-1273 COVID-19 vaccine effectiveness against the SARS-CoV-2 Delta variant in Qatar. *Nat Med*. 2021;27(12):2136–43.
 57. Salman B, Zhou D, Jaffee EM, Edil BH, Zheng L. Vaccine therapy for pancreatic cancer. *Oncoimmunology*. 2013;2(12):e26662.
 58. Lepisto AJ, Moser AJ, Zeh H, Lee K, Bartlett D, McKolanis JR, et al. A phase I/II study of a MUC1 peptide pulsed autologous dendritic cell vaccine as adjuvant therapy in patients with resected pancreatic and biliary tumors. *Cancer Ther*. 2008;6(B):955–64.
 59. Yanagisawa R, Koizumi T, Koya T, Sano K, Koido S, Nagai K, et al. WT1-pulsed dendritic cell vaccine combined with chemotherapy for resected pancreatic cancer in a phase I study. *Anticancer Res*. 2018;38(4):2217–25.
 60. Laheru D, Lutz E, Burke J, Biedrzycki B, Solt S, Onners B, et al. Allogeneic granulocyte macrophage colony-stimulating factor-secreting tumor immunotherapy alone or in sequence with cyclophosphamide for metastatic pancreatic cancer: a pilot study of safety, feasibility, and immune activation. *Clin Cancer Res*. 2008;14(5):1455–63.
 61. Jaffee EM, Hruban RH, Biedrzycki B, Laheru D, Schepers K, Sauter PR, et al. Novel allogeneic granulocyte-macrophage colony-stimulating factor-secreting tumor vaccine for pancreatic cancer: a phase I trial of safety and immune activation. *J Clin Oncol*. 2001;19(1):145–56.
 62. Hardacre JM, Mulcahy M, Small W, Talamonti M, Obel J, Krishnamurthi S, et al. Addition of algenpantucel-L immunotherapy to standard adjuvant therapy for pancreatic cancer: a phase 2 study. *J Gastrointest Surg*. 2012;17(1):94–101.
 63. Romero P, Banchereau J, Bhardwaj N, Cockett M, Disis ML, Dranoff G, et al. The human vaccines project: a roadmap for cancer vaccine development. *Sci Transl Med*. 2016. <https://doi.org/10.1126/scitranslmed.aaf0685>.

64. Keenan BP, Saenger Y, Kafrouni MI, Leubner A, Lauer P, Maitra A, et al. A *Listeria* vaccine and depletion of T-regulatory cells activate immunity against early stage pancreatic intraepithelial neoplasms and prolong survival of mice. *Gastroenterology*. 2014;146(7):1784-94.e6.
65. Yamasaki S, Miura Y, Davydova J, Vickers SM, Yamamoto M. Intravenous genetic mesothelin vaccine based on human adenovirus 40 inhibits growth and metastasis of pancreatic cancer. *Int J Cancer*. 2013;133(1):88-97.
66. Kaufman HL, Kim-Schulze S, Manson K, DeRaffele G, Mitcham J, Seo KS, et al. Poxvirus-based vaccine therapy for patients with advanced pancreatic cancer. *J Transl Med*. 2007;5:60.
67. Hassan R, Thomas A, Alewine C, Le DT, Jaffee EM, Pastan I. Mesothelin immunotherapy for cancer: ready for prime time? *J Clin Oncol*. 2016;34(34):4171-9.
68. Naseri M, Bozorgmehr M, Zöller M, Ranaei Pirmardan E, Madjid Z. Tumor-derived exosomes: the next generation of promising cell-free vaccines in cancer immunotherapy. *Oncoimmunology*. 2020;9(1):1779991.
69. Whiteside TL. The effect of tumor-derived exosomes on immune regulation and cancer immunotherapy. *Future Oncol*. 2017;13(28):2583-92.
70. Xiao L, Erb U, Zhao K, Hackert T, Zoller M. Efficacy of vaccination with tumor-exosome loaded dendritic cells combined with cytotoxic drug treatment in pancreatic cancer. *Oncoimmunology*. 2017;6(6):e1319044.
71. Sun W, Ren Y, Lu Z, Zhao X. The potential roles of exosomes in pancreatic cancer initiation and metastasis. *Mol Cancer*. 2020;19(1):135.
72. Guay C, Krut JK, Rome S, Menoud V, Mulder NL, Jurdzinski A, et al. Lymphocyte-derived exosomal microRNAs promote pancreatic β cell death and may contribute to type 1 diabetes development. *Cell Metab*. 2019;29(2):348-61.e6.
73. Fathollahi A, Hashemi SM, Haji Molla Hoseini M, Yeganeh F. *In vitro* analysis of immunomodulatory effects of mesenchymal stem cell- and tumor cell -derived exosomes on recall antigen-specific responses. *Int Immunopharmacol*. 2019;67:302-10.
74. Lee ES, Sul JH, Shin JM, Shin S, Lee JA, Kim HK, et al. Reactive oxygen species-responsive dendritic cell-derived exosomes for rheumatoid arthritis. *Acta Biomater*. 2021;128:462-73.
75. Maki RG, Livingston PO, Lewis JJ, Janetzki S, Klimstra D, Desantis D, et al. A phase I pilot study of autologous heat shock protein vaccine HSPPC-96 in patients with resected pancreatic adenocarcinoma. *Dig Dis Sci*. 2007;52(8):1964-72.
76. Bloch O, Crane CA, Fuks Y, Kaur R, Aghi MK, Berger MS, et al. Heat-shock protein peptide complex-96 vaccination for recurrent glioblastoma: a phase II, single-arm trial. *Neuro Oncol*. 2014;16(2):274-9.
77. Ji N, Zhang Y, Liu Y, Xie J, Wang Y, Hao S, et al. Heat shock protein peptide complex-96 vaccination for newly diagnosed glioblastoma: a phase I, single-arm trial. *JCI Insight*. 2018;3(10):e99145.
78. Zhang Y, Mudgal P, Wang L, Wu H, Huang N, Alexander PB, et al. T cell receptor repertoire as a prognosis marker for heat shock protein peptide complex-96 vaccine trial against newly diagnosed glioblastoma. *Oncoimmunology*. 2020;9(1):1749476.
79. Wood C, Srivastava P, Bukowski R, Lacombe L, Gorelov AI, Gorelov S, et al. An adjuvant autologous therapeutic vaccine (HSPPC-96; vitespen) versus observation alone for patients at high risk of recurrence after nephrectomy for renal cell carcinoma: a multicentre, open-label, randomised phase III trial. *Lancet*. 2008;372(9633):145-54.
80. Tosti G, Di Pietro A, Ferrucci PF, Testori A. HSPPC-96 vaccine in metastatic melanoma patients: from the state of the art to a possible future. *Expert Rev Vaccines*. 2009;8(11):1513-26.
81. Matsui H, Hazama S, Shindo Y, Nagano H. Combination treatment of advanced pancreatic cancer using novel vaccine and traditional therapies. *Expert Rev Anticancer Ther*. 2018;18(12):1205-17.
82. Gjertsen MK, Bakka A, Breivik J, Saeterdal I, Gedde-Dahl T, Stokke KT, et al. *Ex vivo* ras peptide vaccination in patients with advanced pancreatic cancer: results of a phase I/II study. *Int J Cancer*. 1996;65(4):450-3.
83. Gjertsen MK, Buanes T, Rosseland AR, Bakka A, Gladhaug I, Søreide O, et al. Intradermal ras peptide vaccination with granulocyte-macrophage colony-stimulating factor as adjuvant: clinical and immunological responses in patients with pancreatic adenocarcinoma. *Int J Cancer*. 2001;92(3):441-50.
84. Bernhardt SL, Gjertsen MK, Trachsel S, Møller M, Eriksen JA, Meo M, et al. Telomerase peptide vaccination of patients with non-resectable pancreatic cancer: a dose escalating phase I/II study. *Br J Cancer*. 2006;95(11):1474-82.
85. Gunturu KS, Rossi GR, Saif MW. Immunotherapy updates in pancreatic cancer: are we there yet? *Ther Adv Med Oncol*. 2013;5(1):81-9.
86. Miyazawa M, Ohsawa R, Tsunoda T, Hirono S, Kawai M, Tani M, et al. Phase I clinical trial using peptide vaccine for human vascular endothelial growth factor receptor 2 in combination with gemcitabine for patients with advanced pancreatic cancer. *Cancer Sci*. 2010;101(2):433-9.
87. Suzuki N, Hazama S, Ueno T, Matsui H, Shindo Y, Iida M, et al. A phase I clinical trial of vaccination with KIF20A-derived peptide in combination with gemcitabine for patients with advanced pancreatic cancer. *J Immunother*. 2014;37(1):36-42.
88. Suzuki N, Hazama S, Iguchi H, Uesugi K, Tanaka H, Hirakawa K, et al. Phase II clinical trial of peptide cocktail therapy for patients with advanced pancreatic cancer: VENUS-PC study. *Cancer Sci*. 2017;108(1):73-80.
89. Nishida S, Ishikawa T, Egawa S, Koido S, Yanagimoto H, Ishii J, et al. Combination gemcitabine and WT1 peptide vaccination improves progression-free survival in advanced pancreatic ductal adenocarcinoma: a phase II randomized study. *Cancer Immunol Res*. 2018;6(3):320-31.
90. Gilliam AD, Broome P, Topuzov EG, Garin AM, Pulay I, Humphreys J, et al. An international multicenter randomized controlled trial of G17DT in patients with pancreatic cancer. *Pancreas*. 2012;41(3):374-9.
91. Rong Y, Jin D, Wu W, Lou W, Wang D, Kuang T, et al. Induction of protective and therapeutic anti-pancreatic cancer immunity using a reconstructed MUC1 DNA vaccine. *BMC Cancer*. 2009;9:191.
92. Cappello P, Rolla S, Chiarle R, Principe M, Cavallo F, Perconti G, et al. Vaccination with ENO1 DNA prolongs survival of genetically engineered mice with pancreatic cancer. *Gastroenterology*. 2013;144(5):1098-106.
93. Cappello P, Curcio C, Mandili G, Roux C, Bulfamante S, Novelli F. Next generation immunotherapy for pancreatic cancer: DNA vaccination is seeking new combo partners. *Cancers*. 2018;10(2):51.
94. Zhu K, Qin H, Cha SC, Neelapu SS, Overwijk W, Lizee GA, et al. Survivin DNA vaccine generated specific antitumor effects in pancreatic carcinoma and lymphoma mouse models. *Vaccine*.

- 2007;25(46):7955–61.
95. Niethammer AG, Lubenau H, Mikus G, Knebel P, Hohmann N, Leowardi C, et al. Double-blind, placebo-controlled first in human study to investigate an oral vaccine aimed to elicit an immune reaction against the VEGF-Receptor 2 in patients with stage IV and locally advanced pancreatic cancer. *BMC Cancer*. 2012;12:361.
 96. Fiedler K, Lazzaro S, Lutz J, Rauch S, Heidenreich R. mRNA cancer vaccines. *Recent Results Cancer Res*. 2016;209:61–85.
 97. Wang Y, Zhang Z, Luo J, Han X, Wei Y, Wei X. mRNA vaccine: a potential therapeutic strategy. *Mol Cancer*. 2021;20(1):33.
 98. Pardi N, Hogan MJ, Porter FW, Weissman D. mRNA vaccines - a new era in vaccinology. *Nat Rev Drug Discov*. 2018;17(4):261–79.
 99. Grunwitz C, Kranz LM. mRNA cancer vaccines-messages that prevail. *Curr Top Microbiol Immunol*. 2017;405:145–64.
 100. Alexopoulou L, Holt AC, Medzhitov R, Flavell RA. Recognition of doublestranded RNA and activation of NF-kappaB by Toll-like receptor 3. *Nature*. 2001;413(6857):732–8.
 101. Veetil AT, Zou J, Henderson KW, Jani MS, Shaik SM, Sisodia SS, et al. DNA-based fluorescent probes of NOS2 activity in live brains. *Proc Natl Acad Sci U S A*. 2020;117(26):14694–702.
 102. Heil F, Hemmi H, Hochrein H, Ampenberger F, Kirschning C, Akira S, et al. Species-specific recognition of single-stranded RNA via Toll-like receptor 7 and 8. *Science*. 2004;303(5663):1526–9.
 103. Tanji H, Ohto U, Shibata T, Taoka M, Yamauchi Y, Isoe T, et al. Toll-like receptor 8 senses degradation products of single-stranded RNA. *Nat Struct Mol Biol*. 2015;22(2):109–15.
 104. Neefjes J, Jongmsa MLM, Paul P, Bakke O. Towards a systems understanding of MHC class I and MHC class II antigen presentation. *Nat Rev Immunol*. 2011;11(12):823–36.
 105. Pishesha N, Harmand TJ, Ploegh HL. A guide to antigen processing and presentation. *Nat Rev Immunol*. 2022. <https://doi.org/10.1038/s41577-022-00707-2>.
 106. Roche PA, Furuta K. The ins and outs of MHC class II-mediated antigen processing and presentation. *Nat Rev Immunol*. 2015;15(4):203–16.
 107. Kranz LM, Diken M, Haas H, Kreiter S, Loquai C, Reuter KC, et al. Systemic RNA delivery to dendritic cells exploits antiviral defence for cancer immunotherapy. *Nature*. 2016;534(7607):396–401.
 108. Hasgur S, Yamamoto Y, Fan R, Nicosia M, Gorbacheva V, Zwick D, et al. Macrophage-inducible C-type lectin activates B cells to promote T cell reconstitution in heart allograft recipients. *Am J Transplant*. 2022;22(7):1779–90.
 109. Wykes M, Pombo A, Jenkins C, MacPherson GG. Dendritic cells interact directly with naive B lymphocytes to transfer antigen and initiate class switching in a primary T-dependent response. *J Immunol*. 1998;161(3):1313–9.
 110. Sahin U, Oehm P, Derhovanessian E, Jabulowsky RA, Vormehr M, Gold M, et al. An RNA vaccine drives immunity in checkpoint-inhibitortreated melanoma. *Nature*. 2020;585(7823):107–12.
 111. Kauffman KJ, Webber MJ, Anderson DG. Materials for non-viral intracellular delivery of messenger RNA therapeutics. *J Control Release*. 2016;240:227–34.
 112. Karikó K, Muramatsu H, Welsh FA, Ludwig J, Kato H, Akira S, et al. Incorporation of pseudouridine into mRNA yields superior nonimmunogenic vector with increased translational capacity and biological stability. *Mol Ther*. 2008;16(11):1833–40.
 113. Tsui NBY, Ng EKO, Lo YMD. Stability of endogenous and added RNA in blood specimens, serum, and plasma. *Clin Chem*. 2002;48(10):1647–53.
 114. Weide B, Carralot JP, Reese A, Scheel B, Eigentler TK, Hoerr I, et al. Results of the first phase I/II clinical vaccination trial with direct injection of mRNA. *J Immunother*. 2008;31(2):180–8.
 115. Beck JD, Reidenbach D, Salomon N, Sahin U, Tureci O, Vormehr M, et al. mRNA therapeutics in cancer immunotherapy. *Mol Cancer*. 2021;20(1):69.
 116. Heine A, Juranek S, Brossart P. Clinical and immunological effects of mRNA vaccines in malignant diseases. *Mol Cancer*. 2021;20(1):52.
 117. Weide B, Pascolo S, Scheel B, Derhovanessian E, Pflugfelder A, Eigentler TK, et al. Direct injection of protamine-protected mRNA: results of a phase 1/2 vaccination trial in metastatic melanoma patients. *J Immunother*. 2009;32(5):498–507.
 118. Jackson NAC, Kester KE, Casimiro D, Gurnathan S, DeRosa F. The promise of mRNA vaccines: a biotech and industrial perspective. *NPJ Vaccines*. 2020;5:11.
 119. Pardi N, Muramatsu H, Weissman D, Kariko K. *In vitro* transcription of long RNA containing modified nucleosides. *Methods Mol Biol*. 2013;969:29–42.
 120. Naik R, Peden K. Regulatory considerations on the development of mRNA vaccines. *Curr Top Microbiol Immunol*. 2020. https://doi.org/10.1007/82_2020_220.
 121. Wang RF, Rosenberg SA. Human tumor antigens for cancer vaccine development. *Immunol Rev*. 1999;170(1):85–100.
 122. Leko V, Rosenberg SA. Identifying and targeting human tumor antigens for T cell-based immunotherapy of solid tumors. *Cancer Cell*. 2020;38(4):454–72.
 123. Nejo T, Yamamichi A, Almeida ND, Goretsky YE, Okada H. Tumor antigens in glioma. *Semin Immunol*. 2020;47: 101385.
 124. Li WH, Li YM. Chemical strategies to boost cancer vaccines. *Chem Rev*. 2020;120(20):11420–78.
 125. Lang F, Schrörs B, Löwer M, Tureci O, Sahin U. Identification of neoantigens for individualized therapeutic cancer vaccines. *Nat Rev Drug Discov*. 2022;21(4):261–82.
 126. Abelin JG, Keskin DB, Sarkizova S, Hartigan CR, Zhang W, Sidney J, et al. Mass spectrometry profiling of HLA-associated peptidomes in mono-allelic cells enables more accurate epitope prediction. *Immunity*. 2017;46(2):315–26.
 127. Alspach E, Lussier DM, Miceli AP, Kizhvatov I, DuPage M, Luoma AM, et al. MHC-II neoantigens shape tumour immunity and response to immunotherapy. *Nature*. 2019;574(7780):696–701.
 128. Jurtz V, Paul S, Andreatta M, Marcatili P, Peters B, Nielsen M. NetMHCpan-4.0: improved peptide-MHC class I interaction predictions integrating eluted ligand and peptide binding affinity data. *J Immunol*. 2017;199(9):3360–8.
 129. O'Donnell TJ, Rubinsteyn A, Bonsack M, Riemer AB, Laserson U, Hammerbacher J. MHCflurry: open-source class I MHC binding affinity prediction. *Cell Syst*. 2018;7(1):129–32.e4.
 130. Zhao W, Sher X. Systematically benchmarking peptide-MHC binding predictors: from synthetic to naturally processed epitopes. *PLoS Comput Biol*. 2018;14(11):e1006457.
 131. Blaha DT, Anderson SD, Yoakum DM, Hager MV, Zha Y, Gajewski TF, et al. High-throughput stability screening of neoantigen/HLA complexes improves immunogenicity predictions. *Cancer Immunol Res*. 2019;7(1):50–61.
 132. Rasmussen M, Fenoy E, Harndahl M, Kristensen AB, Nielsen IK, Nielsen M, et al. Pan-specific prediction of peptide-MHC class I complex stability, a correlate of T cell immunogenicity. *J Immunol*. 2016;197(4):1517–24.
 133. Trolle T, Nielsen M. NetTepi: an integrated method for the prediction of T cell epitopes. *Immunogenetics*. 2014;66(7–8):449–

- 56.
134. Weng Y, Li C, Yang T, Hu B, Zhang M, Guo S, et al. The challenge and prospect of mRNA therapeutics landscape. *Biotechnol Adv*. 2020;40:107534.
135. Linares- Fernández S, Lacroix C, Exposito JY, Verrier B. Tailoring mRNA vaccine to balance innate/adaptive immune response. *Trends Mol Med*. 2020;26(3):311–23.
136. Diken M, Kreiter S, Selmi A, Britten CM, Huber C, Türeci Ö, et al. Selective uptake of naked vaccine RNA by dendritic cells is driven by macropinocytosis and abrogated upon DC maturation. *Gene Ther*. 2011;18(7):702–8.
137. Miao L, Zhang Y, Huang L. mRNA vaccine for cancer immunotherapy. *Mol Cancer*. 2021;20(1):41.
138. Victor MP, Acharya D, Begum T, Ghosh TC. The optimization of mRNA expression level by its intrinsic properties—Insights from codon usage pattern and structural stability of mRNA. *Genomics*. 2019;111(6):1292–7.
139. De Haro C, Méndez R, Santoyo J. The eIF-2 α kinases and the control of protein synthesis 1. *FASEB J*. 1996;10(12):1378–87.
140. Liang SL, Quirk D, Zhou A. RNase L: Its biological roles and regulation. *IUBMB Life*. 2006;58(9):508–14.
141. Batey RT, Kieft JS. Improved native affinity purification of RNA. *RNA*. 2007;13(8):1384–9.
142. Karikó K, Muramatsu H, Ludwig J, Weissman D. Generating the optimal mRNA for therapy: HPLC purification eliminates immune activation and improves translation of nucleoside-modified, proteinencoding mRNA. *Nucleic Acids Res*. 2011;39(21):e142.
143. Thess A, Grund S, Mui BL, Hope MJ, Baumhof P, Fotin-Mleczek M, et al. Sequence-engineered mRNA without chemical nucleoside modifications enables an effective protein therapy in large animals. *Mol Ther*. 2015;23(9):1456–64.
144. Weissman D, Karikó K. mRNA: fulfilling the promise of gene therapy. *Mol Ther*. 2015;23(9):1416–7.
145. Mauro VP. Codon optimization in the production of recombinant biotherapeutics: potential risks and considerations. *BioDrugs*. 2018;32(1):69–81.
146. Morrison AH, Byrne KT, Vonderheide RH. Immunotherapy and prevention of pancreatic cancer. *Trends Cancer*. 2018;4(6):418–28.
147. Ho WJ, Jaffee EM, Zheng L. The tumour microenvironment in pancreatic cancer — clinical challenges and opportunities. *Nat Rev Clin Oncol*. 2020;17(9):527–40.
148. Hosein AN, Brekken RA, Maitra A. Pancreatic cancer stroma: an update on therapeutic targeting strategies. *Nat Rev Gastroenterol Hepatol*. 2020;17(8):487–505.
149. Danilova L, Ho WJ, Zhu Q, Vithayathil T, De Jesus-Acosta A, Azad NS, et al. Programmed cell death ligand-1 (PD-L1) and CD8 expression profiling identify an immunologic subtype of pancreatic ductal adenocarcinomas with favorable survival. *Cancer Immunol Res*. 2019;7(6):886–95.
150. Wartenberg M, Cibir S, Zlobec I, Vassella E, Eppenberger-Castori S, Terracciano L, et al. Integrated genomic and immunophenotypic classification of pancreatic cancer reveals three distinct subtypes with prognostic/predictive significance. *Clin Cancer Res*. 2018;24(18):4444–54.
151. Collisson EA, Sadanandam A, Olson P, Gibb WJ, Truitt M, Gu S, et al. Subtypes of pancreatic ductal adenocarcinoma and their differing responses to therapy. *Nat Med*. 2011;17(4):500–3.
152. Waddell N, Pajic M, Patch AM, Chang DK, Kassahn KS, Bailey P, et al. Whole genomes redefine the mutational landscape of pancreatic cancer. *Nature*. 2015;518(7540):495–501.
153. Bailey P, Chang DK, Nones K, Johns AL, Patch AM, Gingras MC, et al. Genomic analyses identify molecular subtypes of pancreatic cancer. *Nature*. 2016;531(7592):47–52.
154. Sivakumar S, de Santiago I, Chlon L, Markowitz F. Master regulators of oncogenic KRAS response in pancreatic cancer: an integrative network biology analysis. *PLoS Med*. 2017;14(1):e1002223.
155. Knudsen ES, Vail P, Balaji U, Ngo H, Botros IW, Makarov V, et al. Stratification of pancreatic ductal adenocarcinoma: combinatorial genetic, stromal, and immunologic markers. *Clin Cancer Res*. 2017;23(15):4429–40.
156. Puleo F, Nicolle R, Blum Y, Cros J, Marisa L, Demetter P, et al. Stratification of pancreatic ductal adenocarcinomas based on tumor and microenvironment features. *Gastroenterology*. 2018;155(6):1999–2013.e3.
157. Karasinska JM, Topham JT, Kalloger SE, Jang GH, Denroche RE, Culibrk L, et al. Altered gene expression along the glycolysis-cholesterol synthesis axis is associated with outcome in pancreatic cancer. *Clin Cancer Res*. 2020;26(1):135–46.
158. Kalimuthu SN, Wilson GW, Grant RC, Seto M, O’Kane G, Vajpeyi R, et al. Morphological classification of pancreatic ductal adenocarcinoma that predicts molecular subtypes and correlates with clinical outcome. *Gut*. 2020;69(2):317–28.
159. Law HCH, Lagundžin D, Clement EJ, Qiao F, Wagner ZS, Krieger KL, et al. The proteomic landscape of pancreatic ductal adenocarcinoma liver metastases identifies molecular subtypes and associations with clinical response. *Clin Cancer Res*. 2020;26(5):1065–76.
160. Murphy JE, Wo JY, Ryan DP, Clark JW, Jiang W, Yeap BY, et al. Total neoadjuvant therapy with FOLFIRINOX in combination with losartan followed by chemoradiotherapy for locally advanced pancreatic cancer: a phase 2 clinical trial. *JAMA Oncol*. 2019;5(7):1020–7.
161. Hackert T, Sachsenmaier M, Hinz U, Schneider L, Michalski CW, Springfield C, et al. Locally advanced pancreatic cancer: neoadjuvant therapy with folfirinox results in resectability in 60% of the patients. *Ann Surg*. 2016;264(3):457–63.
162. Michelakos T, Pergolini I, Castillo CF, Honselmann KC, Cai L, Deshpande V, et al. Predictors of resectability and survival in patients with borderline and locally advanced pancreatic cancer who underwent neoadjuvant treatment with FOLFIRINOX. *Ann Surg*. 2019;269(4):733–40.
163. Singhi AD, Wood LD. Early detection of pancreatic cancer using DNA-based molecular approaches. *Nat Rev Gastroenterol Hepatol*. 2021;18(7):457–68.
164. Fischer CG, Wood LD. From somatic mutation to early detection: insights from molecular characterization of pancreatic cancer precursor lesions. *J Pathol*. 2018;246(4):395–404.
165. Huang X, Zhang G, Liang T. Pancreatic tumor initiation: the potential role of IL-33. *Signal Transduct Target Ther*. 2021;6(1):204.
166. Alonso-Curbelo D, Ho YJ, Burdziak C, Maag JLV, Morris JP, Chandwani R, et al. A gene-environment-induced epigenetic program initiates tumorigenesis. *Nature*. 2021;590(7847):642–8.

<https://doi.org/10.1186/s40779-022-00416-w>

Cite this article as: Huang X, Zhang G, Tang TY, Gao X, Liang TB. Personalized pancreatic cancer therapy: from the perspective of mRNA vaccine. *Mil Med Res*. 2022;9(1):53.