

Physical exercise reverses immuno-cold tumor microenvironment *via* inhibiting SQLE in non-small cell lung cancer

Zhi-Wen Luo^{1†}, Ya-Ying Sun^{1,2†}, Wei Xia^{3†}, Jun-Ying Xu⁴, Dong-Jing Xie⁴, Chun-Meng Jiao⁵, Ji-Ze Dong², Hui Chen⁶, Ren-Wen Wan¹, Shi-Yi Chen^{1*}, Jie Mei^{4*} and Wen-Jun Mao^{7*}

Key words Physical exercise, Non-small cell lung cancer (NSCLC), Squalene epoxidase (SQLE), Tumor immune microenvironment (TIME)

Dear Editor,

Physical exercise has been shown to be associated with reduced cancer incidence and cancer-associated mortality[1,2], but the underlying mechanisms are obscure. Immunometabolic regulation has emerged as one of the most prominent mechanisms explaining the effects of exercise on cancer[1,2]. Physical exercise primarily lowers blood cholesterol and triglycerides, and protects against cardiovascular diseases[3]. However, whether physical exercise can modulate cholesterol metabolism in tumor cells is currently unknown.

Metabolic reprogramming is one of the hallmarks of cancer, and metabolic dysregulation critically contributes toward oncogenesis and tumor progression[4]. Being a common phenomenon associated with both physical exercise and cancer, metabolic regulation is one of the critical mechanisms that mediates the anticancer effects of physical exercise.

Cholesterol, the major sterol in mammalian cell membranes, maintains cell integrity and intracellular homeostasis. Previously, we found that certain cholesterol-related genes were more active in non-small cell lung cancer (NSCLC), which is an immuno-cold tumor. Blocking cholesterol production by treating these cancer cells with 3-hydroxy-3-methylglutaryl-coenzyme A reductase (HMGCR) inhibitors induced an elevated immune response to the inflamed tumor immune microenvironment (TIME)[5]. However, the complexity

of cholesterol biosynthesis warrants the discovery of more interventions.

In this study, we first explored the effects of physical exercise on gene expression in tumors. Re-analyzing the GSE62628 dataset, we found that physical exercise significantly modulated the transcriptome of mouse melanoma cells (Fig. 1a, Additional file 1: Fig. S1). The upregulated genes were associated with immune-related processes (Additional file 1: Fig. S2a, b), and the down-regulated genes were associated with cholesterol biosynthesis (Additional file 1: Fig. S2c, d). We further validated this result using a mouse lung cancer model (Fig. 1b), and found that exercise notably inhibited tumor growth (Fig. 1c, Additional file 1: Fig. S3). Next, we used mass cytometry to analyze changes in the TIME (Additional file 1: Fig. S4). Exercise triggered the infiltration of antitumor immune cells, such as CD8⁺ T cells, M1 macrophages, and B cells, while inhibiting the infiltration of pro-tumor immune cells, such as myeloid-derived suppressor cells (Fig. 1d, Additional file 1: Fig. S5). T cell subset analysis revealed that the proportions of naïve and activated T cells were significantly increased, further evidencing that exercise modulated the TIME (Additional file 1: Fig. S6). The increased proportions of CD8⁺ T cells and M1 macrophages were verified using immunofluorescence analysis of mouse tumor tissues (Fig. 1e). We also validated the effect of exercise on cholesterol metabolic reprogramming in mouse tumor tissues. Exercise significantly inhibited squalene epoxidase (SQLE) expression, but did not affect HMGCR expression (Fig. 1e). Overall, we found that physical exercise reversed the immuno-cold TIME and inhibited cholesterol metabolism.

To identify the critical gene controlling cholesterol metabolism in NSCLC, we investigated the expression and

[†]Zhi-Wen Luo, Ya-Ying Sun, and Wei Xia contributed equally to this work

*Correspondence: cshiyi@163.com; meijie1996@njmu.edu.cn; maowenjun1@njmu.edu.cn

¹Department of Sports Medicine, Huashan Hospital Affiliated to Fudan University, Shanghai 200040, China

²Department of Oncology, the Affiliated Wuxi People's Hospital of Nanjing Medical University, Wuxi 214023, Jiangsu, China

³Department of Cardiothoracic Surgery, the Affiliated Wuxi People's Hospital of Nanjing Medical University, Wuxi 214023, Jiangsu, China

Full list of author information is available at the end of the article

prognostic value of cholesterol biosynthesis-related genes in NSCLC using The Cancer Genome Atlas (TCGA) dataset. Most genes were dysregulated in NSCLC (Additional file 1: Fig. S7a), but only the upregulation of *SQLE* was associated with poor prognosis (Additional file 1: Fig. S7b). Given its critical role in catalyzing the primary oxygenation step of sterol biosynthesis and its prognostic significance, we identified *SQLE* as a potential rate-limiting enzyme in cholesterol production that warranted further analysis. Immunohistochemistry (IHC) analysis of human NSCLC tissues was performed to further validate *SQLE* expression in NSCLC, and the result confirmed that *SQLE* was notably up-regulated (Fig. 1f, g). These findings revealed that *SQLE* was a potential oncogene in NSCLC.

Next, we assessed the immuno-correlation of *SQLE* in NSCLC using the TCGA dataset. High *SQLE* expression was related to the downregulation of most immunomodulators (Additional file 1: Fig. S8a, b). In addition, *SQLE* expression was positively correlated with tumor purity and negatively correlated with immune cell infiltration (Additional file 1: Fig. S8c, d). We also validated the negative correlation between *SQLE* expression and CD8⁺ T cell abundance using an in-house NSCLC cohort (Fig. 1h, i). However, we only observed the correlations between *SQLE* expression and the immuno-cold TIME; the effects of *SQLE* on TIME functionality are still unknown. Furthermore, the Single Cell Expression Atlas of human NSCLC tumors uncovered that *SQLE* was enriched in tumor cells (Fig. 1j, Additional file 1: Fig. S9). Given its negative immuno-correlation in NSCLC, we speculated that *SQLE* may be associated with immunotherapeutic responses. In a combined public cohort, we found that *SQLE* was down-regulated in cases with a poor response, and this was also validated in our recruited NSCLC cohort receiving immune checkpoint inhibitors therapy (Fig. 1k, l; Additional file 1: Fig. S10). Moreover, *SQLE* overexpression notably reversed exercise-mediated tumor inhibition *in vivo* (Fig. 1m-o, Additional file 1: Fig. S11). Overall, *SQLE* expression was related to the immuno-cold TIME and reversed physical exercise-induced tumor inhibition and TIME activation.

In summary, physical exercise inhibited tumor progression by significantly downregulating *SQLE*, which modulated the inflamed TIME and enhanced immune checkpoint inhibitors therapy. In addition, *SQLE* expression was related to poor prognosis and the immunocold TIME (Fig. 1p). Overall, we have clarified the important role of *SQLE* in maintaining the immuno-cold phenotype in NSCLC, and propose physical exercise as an intervention for *SQLE*. However, these observations were made in mice; prospective clinical

trials in humans are warranted before we decide to sensitize immunotherapy using exercise therapy.

Abbreviations

HMGCR: 3-Hydroxy-3-methylglutaryl-coenzyme A reductase; IHC: Immunohistochemistry; NSCLC: Non-small cell lung cancer; TIME: Tumor immune microenvironment; *SQLE*: Squalene epoxidase; TCGA: The Cancer Genome Atlas.

Supplementary information

The online version contains supplementary material available at <https://doi.org/10.1186/s40779-023-00474-8>.

Additional file 1. Materials and methods. **Fig. S1** A total of 1980 genes were downregulated and 1445 genes were upregulated. **Fig. S2** Reactome analysis of DEGs in the GSE62628 dataset. **Fig. S3** Weight of the harvested tumors in tumor-bearing mice with or without exercise ($n=5$ per group). **Fig. S4** t-SNE plots of normalized marker expression for classifying tumor cells from mice in the control and exercise groups. **Fig. S5** Marker expression and cell proportion of different cell subpopulations. **Fig. S6** Changes in the T cell subset proportions in mice tumors from the control and exercise groups. **Fig. S7** Expression and prognostic value of genes related to cholesterol synthesis in NSCLC tissues. **Fig. S8** Correlation between *SQLE* expression and TIME features. **Fig. S9** Expression levels of *SQLE* in different cell types in NSCLC tissues in the GSE117570 dataset. **Fig. S10** Representative images depicting *SQLE* expression in patients with different immunotherapy responses and semi-quantitative analysis of the predictive value of *SQLE* ($n=14$ in the non-responder group, $n=8$ in the responder group). **Fig. S11** Weight of the harvested tumors from tumor-bearing mice ($n=5$ per group). **Table S1** A list of the antibodies and reagents used in this research.

Acknowledgements

Not applicable.

Authors' contributions

WJM, JM and SYC contributed to the conception of the study. ZWL, YYS and WX invented the materials and methods. ZWL, YYS and WX carried out the mechanical tests and analysis. ZWL, YYS, WX, CMJ, JZD, HC and RWW designed and conducted the *in vitro* and *in vivo* experiments. JYX and DJX collected the clinical samples. CMJ and JZD helped visualize the data. ZWL, YYS and WX wrote the original manuscript. WJM, JM and SYC reviewed and edited the manuscript. WJM, JM and SYC supervised the project. All authors read and approved the final manuscript.

Funding

This work was supported by the National Natural Science Foundation of China (82172511), the Natural Science Foundation of Jiangsu Province (BK20210068), the Sanming Project of Medicine in Shenzhen (SZSM201612078), the Health Shanghai Initiative Special Fund [Medical-Sports Integration (JKSHZX-2022-02)], the Top Talent Support Program for Young- and Middle-aged People of Wuxi Municipal Health Commission (HB2020003), the Mega-project of Wuxi Commission of Health (Z202216), and the High-end Medical Expert Team of the 2019 Taihu Talent Plan (2019-THRCTD-1).

Availability of data and materials

All data supporting the results of this study are shown in this




Fig. 1 (See legend on next page.)

(See figure on previous page.)

Fig. 1 Physical exercise “heats” immuno-cold tumors by inhibiting cholesterol synthesis.

a. Overview of DEGs between tumors in mice with or without voluntary wheel running; b. Schematic protocol of the experimental procedures in C57BL/6 mice; c. Tumor growth in Lewis carcinoma-bearing mice with or without exercise ($n=5$); d. t-SNE plots of 50,000 cells per group colored using PhenoGraphcluster; e. Immunofluorescence revealing the infiltration of CD8⁺ T cells and M1 macrophages, and the expression of HMGCR and SQLE. Representative images demonstrating SQLE expression in para-tumor and tumor tissues using anti-SQLE staining (f) and semi-quantitative analysis (g); h. Representative images showing CD8 expression in the high- and low-SQLE groups; i. Semi-quantitative analysis of the correlation between SQLE and CD8 expression; j. Expression levels of SQLE in different cell types from NSCLC tissues in 6 scRNA-seq datasets; k and l. Expression of SQLE in patients with different immunotherapy responses and ROC analysis of the predictive value of SQLE ($n=30$ in the non-responder group, $n=13$ in the responder group). Data obtained by merging the GSE126044 and GSE135222 datasets; m. Tumor growth curve and SQLE overexpression in tumors from mice with exercise ($n=5$); n. Representative images showing the tumors harvested from Lewis carcinoma-bearing mice ($n=5$); o. Representative images showing the infiltration of CD8⁺ T cells and M1 macrophages in the above tumors; p. Schematic diagram of this study: exercise heats up the TIME by suppressing SQLE expression. $**P<0.01$, $***P<0.001$. MDSC. Myeloid-derived suppressor cell; IRS. Immunoreactivity score; DEGs. Differentially expressed genes; HMGCR. 3-hydroxy-3-methylglutaryl-coenzyme A reductase; SQLE. Squalene epoxidase; NSCLC. Non-small cell lung cancer; scRNA-seq. Single-cell RNA sequencing; OE. Overexpression; TIME. Tumor immune microenvironment; ROC. Receiver operating characteristic; t-SNE. t-distributed Stochastic Neighbor Embedding; DPT. Double positive T cells

published article and supplementary documents. In addition, original omics data for bioinformatics analysis could be obtained from corresponding platforms.

Declarations

Ethics approval and consent to participate

Ethical approval for the use of tissue microarrays was granted by the Clinical Research Ethics Committee at Outdo Biotech (YBM-05-02, HLugA150CS01, Shanghai, China). In addition, ethical approval for the collection of samples was granted by Clinical Research Ethics Committee at the Affiliated Wuxi People’s Hospital of Nanjing Medical University (KY21126). All animal experiments were approved by the Laboratory Animal Ethics Committee at Nanjing Medical University (2022370).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Sports Medicine, Huashan Hospital Affiliated to Fudan University, Shanghai 200040, China. ²Department of Sports Medicine, Shanghai General Hospital, Shanghai 200080, China. ³Department of Critical Care Medicine, the Affiliated Wuxi People’s Hospital of Nanjing Medical University, Wuxi 214023, Jiangsu, China. ⁴Department of Oncology, the Affiliated Wuxi People’s Hospital of Nanjing Medical University, Wuxi 214023, Jiangsu, China. ⁵Guangdong Pharmaceutical

University, Guangzhou 510006, China. ⁶Institute of Acupuncture Research, Institutes of Integrative Medicine, Fudan University, Shanghai 200433, China. ⁷Department of Cardiothoracic Surgery, the Affiliated Wuxi People’s Hospital of Nanjing Medical University, Wuxi 214023, Jiangsu, China.

References

1. Kurz E, Hirsch CA, Dalton T, Shadaloey SA, Khodadadi-Jamayran A, Miller G, et al. Exercise-induced engagement of the IL-15/IL-15Ra axis promotes anti-tumor immunity in pancreatic cancer. *Cancer Cell*. 2022;40(7):720-37.e5.
2. Rundqvist H, Velica P, Barbieri L, Gameiro PA, Bargiela D, Gojkovic M, et al. Cytotoxic T-cells mediate exercise-induced reductions in tumor growth. *Elife*. 2020;9:e59996.
3. Muscella A, Stefano E, Marsigliante S. The effects of exercise training on lipid metabolism and coronary heart disease. *Am J Physiol Heart Circ Physiol*. 2020;319(1):H76–88.
4. Gyamfi J, Kim J, Choi J. Cancer as a metabolic disorder. *Int J Mol Sci*. 2022;23(3):1155.
5. Mao W, Cai Y, Chen D, Jiang G, Xu Y, Chen R, et al. Statin shapes inflamed tumor microenvironment and enhances immune checkpoint blockade in non-small cell lung cancer. *JCI Insight*. 2022;7(18):e161940.

<https://doi.org/10.1186/s40779-023-00474-8>

Cite this article as: Luo ZW, Sun YY, Xia W, Xu JY, Xie DJ, Jiao CM, et al. Physical exercise reverses immuno-cold tumor microenvironment via inhibiting SQLE in non-small cell lung cancer. *Mil Med Res*. 2023;10(1):39.